

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07C , C07D , C07J , A61K , A01N</p>	<p>A1</p>	<p>(11) International Publication Number: WO 98/32718 (43) International Publication Date: 30 July 1998 (30.07.98)</p>
<p>(21) International Application Number: PCT/NO98/00021 (22) International Filing Date: 23 January 1998 (23.01.98) (30) Priority Data: 9701441.9 24 January 1997 (24.01.97) GB (71) Applicant (for all designated States except US): NORSK HYDRO ASA [NO/NO]; N-0240 Oslo (NO). (72) Inventors; and (75) Inventors/Applicants (for US only): MYHREN, Finn [NO/NO]; Leirkupgt. 7, N-3915 Porsgrunn (NO). BØR-RETZEN, Bernt [NO/NO]; Granlia 3, N-3940 Heistad (NO). DALEN, Are [NO/NO]; Nordahl Brunsvei 7, N-7015 Trondheim (NO). SANDVOLD, Marit, Liland [NO/NO]; Skiferveien 31, N-3931 Porsgrunn (NO). (74) Agent: LILLEGRAVEN, Rita; Norsk Hydro ASA, N-0240 Oslo (NO).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: NEW FATTY ACID DERIVATIVES (57) Abstract The properties of biologically active compounds, for example drugs and agrochemicals, which contain in their molecular structure one or more functional groups selected from alcohol, ether, phenyl, amino, amido, thiol, carboxylic acid and carboxylic acid ester groups are modified by replacing one or more of these functional groups by a lipophilic group selected from those of the formula: RCOO-, RCONH-, RCOS-, RCH₂O-, RCH₂NH-, -COOCH₂R, -CONHCH₂R and -SCH₂R, wherein R is a lipophilic moiety selected from cis-8-heptadecenyl, trans-8-heptadecenyl, cis-10-nonadecenyl and trans-10-nonadecenyl.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

NEW FATTY ACID DERIVATIVES

This invention relates to biologically active compounds, and it is concerned with providing by means of chemical
5 derivatisation, a technique whereby the behaviour of many biologically active compounds, such as drugs and agricultural chemicals, may be favourably modified.

There is a great interest in the medical community to
10 investigate and improve the transport efficiency of drugs to the site of action in the patient. The work has mainly been focused on resorption of a drug from the intestine to the blood stream, although transport across other biological barriers often plays an important role
15 in obtaining the necessary therapeutic effect in the treatment of many diseases like cancer, infections, inflammations, CNS disorders etc. The transport across the cell membrane is often a main impediment to achieve optimal effect with a therapeutic compound.

20 Over the last decades drug resistance in the treatment of malignant and infectious diseases has become increasingly prevalent and is now regarded as a serious clinical problem. The development of drug resistance can be due
25 to a number of mechanisms, but quite often relates to a triggering of the normal mechanisms whereby microorganisms and cells clear toxic compounds to subtoxic levels. One example is the development of multi-drug resistance (MDR) in cancer cells. In this
30 case MDR frequently relates to a cellular membrane protein pump by which the cells achieve a very efficient efflux of toxic compounds. In a clinical situation, i.e. the treatment of a tumour with a cytostatic drug, cells with the most potent protein pump can
35 preferentially survive, and these cells may proliferate to a new tumour which may be resistant to treatment with a variety of different drugs. Similar mechanisms of action may be responsible for the lack of effect seen in other therapeutic areas, for instance with anti-malarial

drugs.

Several techniques to try to circumvent resistance mechanisms in the clinic are known. For example, the
5 co-administration of a Ca^{2+} channel blocker such as verapamil or an immunomodulating agent like cyclosporin, have been tried out. However, no significant improvements have been reported so far.

10 There have been several proposals in the literature for improving the therapeutic index, bioavailability, membrane passage, organ targeting, etc of therapeutic compounds by combining the compounds with fatty acids so
15 as to form either chemically coupled derivatives or physical mixtures.

Thus, for example, EP-A-393920 discloses that anti-viral nucleosides and nucleoside analogues which are
20 derivatised with long chain (C_{16} upwards) acyl groups have advantages as compared with the parent compound. It is stated that the fatty acid portion of these molecules preferably are made up of polyunsaturated fatty acids, such as γ -linolenic or linoleic acid.

25 US-A-3920630 teaches that 2,2'-anhydro-aracytidine and its 5'-O-acylates have the same general biological and therapeutic activity as anti-viral agents as ara-cytidine itself. The compound 2,2'-anhydro-5'-O-oleyl-ara-cytidine is specially mentioned.

30 EP-A-56265 discloses esters of arabino-furanosyl-thymine (Ara T) with saturated acids having 1-17 C-atoms.

From PCT/WO90/00555 there are known lipid derivatives
35 linked, especially through a phosphate group, to the 5'-position of the pentose group of a nucleoside. The purpose of this derivatisation is to make the nucleosides more lipophilic so that they could be included into

liposomes, which are preferentially taken up by macrophages and monocytes, cells which are found to harbour the HIV virus. It is stated that a targeting effect is thereby achieved.

5

The anti-viral and anti-cancer activities of nucleoside analogues are directly linked to intra-cellular phosphorylation of the administered drug.

10 This biochemical transformation is normally effectuated by viral and/or cellular enzymes. To improve the effect WO96/25421 discloses phospholipid derivatives of nucleosides with relatively short chain (C_{14} or less) saturated or unsaturated fatty acids.

15 The art has also sought to improve the characteristics of other classes of pharmaceutical substance through derivatisation with fatty acids.

For example, WO96/22303 teaches that the pharmokinetic
20 profile and mode of delivery of several different categories of therapeutic compounds (corticosterones, opioids and opioid antagonists, anti-viral nucleosides, cyclosporins and related cyclopeptides, folate antagonists, catecholamine precursors and catecholamines
25 and alkylating agents containing a carboxylic acid group) can be altered by conjugating them to one to three acyl derivatives of fatty acids through the use of a linker/spacer group which includes a tromethamine or ethanolamine derivative. Palmitic acid is the preferred
30 fatty acid.

Lipophilic pro drugs of several NSAIDs are known from H. Bundgaard et al (International Journal of
Pharmaceutics, 43 101-110 1988) and V.R. Shanbhag et al
35 (Journal of Pharmaceutical Sciences, 149 Vol. 81, No. 2, February 1992). In addition to the pro drug aspect, reduced GI irritation is reported. EP-A-0195570 suggests that the administration of gamma-linolenic and

dihomo-gamma-linolenic acid in conjunction with NSAIDs reduces the side effects shown by the NSAIDs when taken on a continuing basis.

- 5 USP 5,284,876 teaches the use of docosahexaenoic acid amides of dopamine as per oral pro drugs in the treatment of CNS disorders.

- Physical mixtures containing fatty acids/fatty acid
10 derivatives used as so-called penetration enhancers both with dermal and per oral administration are known from PCT/US94/02880 and PCT/SE96/00122.

- As indicated, many of these prior proposals concern fatty
15 acid derivatives of anti-viral nucleosides and nucleoside analogues. It is indeed not surprising that this should be so as it has long been known that certain polyunsaturated fatty acids attack viruses.

- In EP-A-0642525 we ourselves taught that the anti-viral
20 effect of nucleosides and nucleoside analogues can be highly potentiated through reaction with oleic acid (cis-9-octadecenoic acid), elaidic acid (trans-9-octadecenoic acid), cis-11-eicosenoic acid or trans-11-eicosenoic acid, to form the corresponding 5'-O-
25 monoester. We have shown that the beneficial effects which can be obtained from these four specific monounsaturated, ω -9 C18 or C20 fatty acids are superior to those generally obtainable through fatty acid derivatisation.

- 30 We have now surprisingly found in accordance with the present invention that the properties of numerous different biologically active compounds may be favourably modified by derivatisation with an ω -9 C18 or C20
35 monounsaturated fatty acid. The present invention thus provides a widely utilisable but simple technique for enhancing the value of many drugs and agricultural chemicals, for instance.

Broadly, the present invention in one aspect provides a lipophilic derivative of a biologically active compound containing in its molecular structure one or more
5 functional groups selected from alcohol, ether, phenyl, amino, amido, thiol, carboxylic acid and carboxylic acid ester groups, other than a nucleoside or nucleoside analogue, said lipophilic derivative being characterised by a molecular structure in which the or at least one
10 said functional group of said biologically active compound is replaced by a lipophilic group selected from those of the formula: RCOO- , RCONH- , RCOS- , $\text{RCH}_2\text{O-}$, $\text{RCH}_2\text{NH-}$, $\text{-COOCH}_2\text{R}$, $\text{-CONHCH}_2\text{R}$ and $\text{-SCH}_2\text{R}$, wherein R is a lipophilic moiety selected from cis-8-heptadecenyl,
15 trans-8-heptadecenyl, cis-10-nonadecenyl and trans-10-nonadecenyl.

In one preferred embodiment of the present invention, the biological effect of a therapeutic compound is improved
20 by derivatising the compound with a n-9 C18 or C20 monounsaturated fatty acid. We present below a detailed discussion of the application of this invention to drugs selected from the following groups:

- 25 1. cancer drugs;
2. antiinflammatory drugs
 - NSAIDs
 - adrenocorticosteroids;
3. antibiotics and other antibacterial agents;
- 30 4. antiparasitic drugs;
5. CNS drugs;
6. cardiovascular drugs; and
7. anticoagulants.

35 However, the invention is broadly applicable to any compound which is pharmacologically active and which possesses in its molecule one or more functional groups capable of conjugating with an n-9 C18 or C20

monounsaturated fatty acid. Thus, for example, the present invention may also be used to improve the biological effects of medicinal compounds of the following types, for instance analgesics, fungicides, antihyperlipidemics, antiemetics and diagnostics.

The lipophilic derivatives of therapeutically active compounds in accordance with the present invention may be formulated with pharmaceutically acceptable carriers and excipients by conventional procedures well known to those skilled in the art. The dosage rates will be correlated with those of the mother drug, although in cases where the lipophilic derivatives of the invention strongly potentiate the effect of the mother drug it may be possible to reduce the dosage from normal levels.

Although the beneficial effects of the present invention have been demonstrated on well-established drugs, it is believed that similar improvements are likely to be exhibited by other drugs which are still in the course of development. That is to say, the postulated explanation for the improved properties which we have observed with lipophilic derivatives of this invention, is of general application and not restricted to any specific mechanism for therapeutic activity.

A particularly valuable property which we have found exhibited by some of the lipophilic derivatives of therapeutically active compounds in accordance with the present invention is that they overcome drug resistance. Although we do not wish to be bound by theory, it is believed that the lipophilic derivatives of this invention interact in some manner with membrane protein pumps so that the cells are inhibited from clearing the active (toxic) compounds, thus enabling the concentration of the active compounds to be sustained at a therapeutically beneficial level for longer periods. In any event, the present invention leads also to the

possibility of combatting the effects of drug resistance by the co-administration of a mother drug and a lipophilic derivative of that drug in accordance with the present invention. Suitably, the mother drug and
5 lipophilic derivative thereof will normally be presented in the same pharmaceutical preparation for ease of administration, although in some cases it may be preferred to present the mother drug and lipophilic derivative in separate unit dosage forms. The dosage of
10 the lipophilic derivative, relative to that of the mother drug can be determined by appropriate tests but generally will range from 1:1 to 1000:1 by weight.

As previously indicated, the invention is generally
15 applicable to any compound having biological activity and not just drugs.

Another economically important class of biologically active compounds are products used in agriculture and
20 horticulture, for example pesticides, fungicides and herbicides. Agrochemicals vary widely, both in structure and in their modes of action. For instance, there are several well-recognized routes of uptake; for example plants may take up the active compound either
25 through the root system or directly through the leaves or stem of the plant, while a pesticide may be taken up either through a plant which the pest attacks or by direct contact. Lipophilic derivatives of agrochemicals in accordance with this invention are found to have
30 enhanced take up potential both by plants and by insects and other pests. Moreover, the present derivatives help to combat pesticide resistance which, like drug resistance, is a growing problem.

35 Other classes of biologically active compounds which can with advantage be derivatised in accordance with the present invention include food and feed additives such as conserving agents, fragrances and spices.

The lipophilic derivatives of the present invention may be prepared by reacting the parent drug or other biologically active compound molecule with a cis- or trans-n-9 monounsaturated fatty acid, fatty acid alcohol or fatty amine having a chain length of 18 or 20 carbon atoms, or with a reactive derivative of such a fatty acid, fatty alcohol or fatty amine, for example acid chlorides, reactive esters, halogenides or the like. The notation n-9 indicates that the unsaturation is between the 9 and 10 positions counted from the C-terminal of the lipidic moiety. Thus, that fatty acids (and alcohols and amines derived therefrom) which may be used are cis-9-octadecenoic acid (oleic acid), trans-9-octadecenoic acid (elaidic acid), cis-11-eicosenoic acid and trans-11-eicosenoic acid.

The coupling reaction between the parent biologically active compound and the fatty acid, fatty alcohol or fatty amine compound can be accomplished by a variety of methods known to those skilled in the art. When two or more derivatisable functional groups are present in the parent molecule, then protecting groups or modified synthetic methods may be used to achieve the necessary selectivity in the coupling steps. In general, the progress of the reactions can be followed using thin layer chromatography (TLC) and appropriate solvent systems. When the reaction is completed as determined by TLC, the product is generally extracted with an organic solvent and purified by chromatography and/or recrystallization from an appropriate solvent system. If more than one hydroxyl, amino, thiol or carboxylic group is present in the parent starting material, a mixture of alkylated or acylated compounds may be produced. The individual mono- or poly-derivatised compounds may then be separated by, for instance, chromatography.

Often, the coupling reaction may be accomplished in one-step, and generally the lipophilic derivatives can be recovered as crystals with good stability profiles, which is helpful for successful galenical processing of the finished pharmaceutical product.

The preparative processes which may be used according to the present invention are illustrated by the reaction schemes given below as well as by the working examples given later in this specification.

The invention will now be further described in detail in connection with several different categories of drugs.

15 Antiinflammatory Drugs

A number of serious diseases, such as rheumatoid arthritis, osteoarthritis, Bechterews syndrome, systemic lupus erythematosus (SLE), asthma, gout, etc are a result of an abnormal immune response eliciting an inflammatory reaction. The inflammatory process involves a series of events that can be elicited by a number of stimuli, for example antigen-antibody interactions, infectious agents, ischemia, etc. At a macroscopic level, the response is usually accompanied by the clinical signs of erythema, edema, tenderness (hyperalgesia) and pain. Inflammatory diseases are mainly treated with three types of drugs, namely NSAIDs (also sometimes termed aspirin-like drugs), immunosuppressive drugs (e.g. methotrexat, cyclophosphamide and lately also cyclosporin) and adrenocorticosteroids (hydrocortisone, prednisolone, etc). The treatment mainly suppresses the pain and/or intensity of the onset of the disease. Current treatment regimens are often limited by severe side effects due to high dosing and/or prolonged treatment periods.

Reversible airway obstruction -asthma- is the most common

of the breathing disorders. The degree of bronchial hyperresponsiveness is normally controlled or reduced by the regular inhalation of adrenocortical steroids and/or bronchodilators. Most treatment regimens provide 2-3
5 hours of therapeutic effect on average. For most respiratory aerosols, absorption is virtually equivalent to that from parenteral or oral administration. Treatment is palliative by virtue of the antiinflammatory and immunosuppressive effects. For prolonged treatment,
10 the smallest dose that will depress side effects is to be used.

Methylprednisolone sodium succinate is given by intravenous administration, followed by oral
15 administration for up to 10 days in serious attacks of asthma. Acute exacerbation of asthma is often treated with brief courses of oral corticosteroids. Incorporation of inhaled corticosteroids in regimens for the treatment of bronchial asthma has increased
20 substantially in recent years. Beclomethasone dipropionate, tramcinolone acetonide or flunisolide can either reduce the duration of courses of oral corticosteroids or replace them entirely. Less suppression of adrenal function is seen when the
25 drugs are used at the recommended doses.

The use of steroids in local treatment of chronic asthma can conveniently be done by means of inhalators. This limits the risk of severe side effects following systemic
30 administration. To assure rapid and selective action following local administration, addition to and interaction with receptors on endothelial cells in the bronchia is essential. The fatty acid derivatives of this invention, with their ability to anchor the active
35 drug to the cells, can further improve the benefit gained from local administration. Inflammation of the airways is recognised as a prominent feature of fatal asthma attacks, and similar changes have been found in bronchial

biopsies of even mild asthma attacks. Severe asthma attacks are associated with a high influx of inflammatory cells in the airways, mainly alveolar macrophages. This situation is quite similar to the appearance of
5 airway hyperresponsiveness induced by viruses. \


Macrophages can release reactive oxygen species which increase pulmonary resistance and induce histamine release. Models that measure the suppression of inflammatory cells in general and macrophages in
10 particular can be used to evaluate the effect of possible asthma drugs. Chemoluminescence can be used as a measure of the release of reactive oxygen species. As shown below, the influx of inflammatory cells into the peritoneum of rats, mainly macrophages is reduced by
15 adrenocorticosteroid derivatives of this invention. The activity of the inflammatory cells upon stimulation is also reduced, and the reduction is observed for a prolonged period of time post treatment for the derivatives compared to their mother drugs.

20 The derivatives are active for at least 48 hours post treatment. This prolonged activity may give a great advantage in asthma treatment.

Natural hormones are often being so rapidly degraded
25 in vivo that unless injected frequently little therapeutic benefit can be achieved by combining these molecules, or synthetic analogues mimicking the natural compounds, with the fatty acids described in this invention, the pharmacokinetic behaviour can be altered
30 to improve the therapeutic benefit. This is valid for both systemic and local administration.


Examples of adrenocorticosteroids and other asthma drugs which may be derivatised in accordance with the present
35 invention include:

12




BETAMETHASONE

5




CORTISONE




DEXAMETHASONE

10




FLUOCINOLONE




15

FLUDROCORTISONE




HYDROCORTISONE




20

METHYLPREDNISOLONE




PARAMETHASONE

13




PREDNISOLONE




PREDNISONE

5




TRIAMCINOLONE




BECLOMETHASONE

10



EPROZINOL



ORCIPRENALIN

15

The most commonly used drugs in the treatment of inflammatory diseases are the NSAIDs. There exist many different products of this class in frequent use, the leading ones being naproxen, diclofenac (voltaren), piroxicam (felden) and salicylic acid derivatives. NSAIDs are antiinflammatory, analgesic and antipyretic but their main clinical uses are in the treatment of inflammatory disorders. NSAIDs mainly provide symptomatic relief from the pain and inflammation associated with these diseases but they do not arrest the progression of the pathological injury of tissue during severe episodes. None of the NSAID products known today reduce the formation of granulomatous tissue to any significant extent. Although a reduction in

granulomatous fluid content is observed, that effect is not reflected in a simultaneous reduction in granulomatous solid matter content. The chief mode of action of these drugs is inhibition of the prostaglandin bio-synthesis (inhibition of cyclooxygenase). The distribution and pharmacokinetic properties of each agent have an important impact on the drug's activity. This is also believed to be the reason for the great variation in the response of individual patients to different NSAID drugs, even those in the same chemical family. For example, large variations in tolerance to the different propionic acid derivatives are reported.


The main property of NSAIDs is their ability to inhibit cyclooxygenase and hence the bio-synthesis of PGG₂ and PGH₂ and all the eicosanoids derived therefrom (PGI₂, TXB₂, PGE₂, etc). On the other hand NSAIDs are not known to inhibit the lipooxygenase (at least not to the same degree) and therefore do not influence the synthesis of the leukotrienes (LTB₄ and LTC₄). The prostaglandins PGI₂ and PGE₂ play an important role in the inflammatory process. They cause edema and probably increase the vascular permeability. PGI₂ is the main factor of the pain associated with inflammatory diseases. The leukotrienes are important mediators in the second and third phases of an inflammatory attack and as the NSAIDs do not inhibit lipooxygenase to a useful therapeutic extent, they do not influence the degenerative part of the inflammatory disease.

NSAIDs are associated with side effects which occasionally can be severe. The most common side effect shown is a propensity to induce gastric or intestinal ulceration, and hence pain, nausea, heartburn and sometimes bleeding and anaemia. These effects are correlated with inhibition of the bio-synthesis of prostaglandins. As a result of the absence of PGI₂ and TXB₂ the blood platelets lose the ability to aggregate


which in turn results in a longer bleeding time. In many cases it is clear that NSAIDs have no beneficial effect on rheumatoid disease progression, and there is evidence to suggest that in some circumstances they could even
5 accelerate the disease process. This is manifested as severe matrix loss due to increased cartilage breakdown.

Other side effects such as retention of salt and water, hyperkalemia and decrease in renal blood flow are also related to the inhibition of prostaglandin synthesis, and
10 can make treatment impossible. Also in some patients hypersensitivity to aspirin, which can lead to anaphylactic shock, excludes treatment with aspirin like drugs.

15 The parent NSAID may be any compound which can be categorized as a non-steroidal antiinflammatory drug and which possesses one or more derivatisable groups selected from alcohol, ether, phenol, amino (primary, secondary or tertiary), amido, thiol, carboxylic acid and carboxylic
20 ester groups. Currently known NSAIDs of this class include the following compounds:




Acemetacin




Alclofenac

25




Amfenac




Aspirin


16




Bendazac




Benorylate




Benoxaprofen




Bucloxic Acid




Bufexamac




Bumadizon




Butibufen



Carprofen



Cinmetacin



Clidanac


5

10


15

20


17




Clometacin




Clopirac




Diclofenac




Diflunisal




Etodolac




Etofenamate




Felbinac



Fenbufen



Fenclofenac




Fenclorac

5


10

15


18




Fendosal



Fenoprofen




Fentiazac




Flufenamic acid

5




Flurbiprofen


10



Glafenine




Ibufenac




Ibuprofen

15




Indomethacin




Isofezolac

19




Isoxepac




Ketoprofen


5



Ketorolac




Lonazolac




10

Meclofenamic Acid




Mefenamic Acid




Metiazinic Acid


15



Nabumetone




Naproxen




Niflumic Acid

20


20




Oxametacin




Oxaprozin




Pirazolac




Piroxicam




Protizinic Acid




Salicylic Acid




Sulindac



Surgam




Tenidap




Tenoxicam

5


10




Tiaramide




Tinoridine



Tolfenamic Acid



Tolmetin



Zomepirac

As illustrated above, a number of known NSAIDs contain more than one derivatisable group of the above-defined kinds. In these cases, one or more of these functional groups can be replaced by a lipophilic group in accordance with the present invention, and where there are two or more lipophilic groups these may be the same or different lipophilic groups.

The lipophilic antiinflammatory drug derivatives of the present invention may be prepared by reacting the parent drug with a cis or trans n-9 monounsaturated fatty acid, fatty alcohol or fatty amine having a chain length of 18 or 20 carbon atoms, or with a reactive derivative of such a fatty acid, fatty alcohol or fatty amine, for example acid chlorides, reactive esters, halogenides and mesylates. The notation n-9 indicates that the

unsaturation is between the 9 and 10 positions counted from the C-terminal of the lipidic moiety. Thus, the fatty acids (and alcohols and amines derived therefrom) which may be used are cis-9-octadecenoic acid (oleic acid), trans-9-octadecenoic acid (elaidic acid), cis-11-eicosenoic acid and trans-11-eicosenoic acid.


The coupling reaction between the parent drug and the fatty acid, fatty alcohol or fatty amine compound can be accomplished by a variety of methods as known to those skilled in the art. When two or more derivatisable functional groups are present in the parent drug, then protecting groups or modified synthetic methods may be used to achieve the necessary selectivity in the coupling steps.

In general, the progress of the reactions can be followed using thin layer chromatography (TLC) and appropriate solvent systems. When the reaction is completed as determined by TLC, the product is generally extracted with an organic solvent and purified by chromatography and/or recrystallization from an appropriate solvent system. If more than one hydroxyl, amino, thiol or carboxylic group is present in the NSAID starting material, a mixture of alkylated or acylated compounds may be produced. The individual mono- or poly-derivatised compounds may then be separated by, for instance, chromatography.

The preparative processes which may be used according to the present invention are illustrated by the reaction schemes given below as well as by the Examples given later in this specification.

The first reaction scheme illustrates the derivatisation of salicylic acid

23




Scheme 1

Treatment of sodium salicylate with a fatty-alcohol
 5 mesylate (R'-OMS) gives the salicylic acid ester (I). A
 modification of this reaction gives the salicylic acid-
 ester-2-ether (II) wherein the hydrocarbon residue in the
 ester and the ether are the same. This product is more
 favourably obtained through the alkylation of ethyl-
 10 salicylate (IV) to give the ethyl salicylate-2-ether (V)
 with a subsequent hydrolysis of the ethyl ester to give
 the salicylic acid-2-ether (III).

A combination of these methods makes possible the product
 15 of di-adducts of formula II but wherein the ester and
 ether substituents are different.

The second reaction scheme illustrates the derivatisation
 of naproxen.

24



Scheme 2.

5 The naproxen ester (VII) or amide (VIII) derivatives are prepared from naproxen (VI) and the corresponding alcohol or amine ($R'-OH$ or $R'-NH_2$) using coupling reagents such as N,N'-dicyclohexylcarbodiimide (DCC) or O-(1H-benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU).


15 The long chain ether analogues (XII) are made from naproxen (VI) utilizing an initial demethylation of the aromatic 6-methylether to give the product (IX) with subsequent esterification of the propionic acid side chain (X). Alkylation of the phenolic moiety (XI) and hydrolysis of the ethyl-ester gave the product (XII). A combination of these methods can give di-adducts

20 wherein the hydrocarbon residue in the ether and the ester or amide are different or the same.

The third reaction scheme illustrates the derivatisation of piroxicam.

25

25




Scheme 3.

5

The piroxicam ester (XIV) is prepared from piroxicam (XIII) and the corresponding fatty acid chloride (R'COCl). Acylation at the amide nitrogen in piroxicam is possible, and a small amount of the N-acylated as well as the di-acylated product is isolated. The identity of the main product (XIV) is confirmed by advanced NMR techniques.

15 The fourth reaction scheme illustrates the derivatisation of diclofenac.




Scheme 4

20

The diclofenac ester (XVI) or amide (VXII) are prepared from diclofenac (XV) and the corresponding alcohol or amine (R-OH or R-NH₂) using coupling reagents such as DCC or TBTU. The isomeric amide (XVIII) can be made from the corresponding fatty acid R'-COOH and XV using TBTU as a coupling reagent.

The fifth reaction scheme illustrates the derivatisation of betamethasone (XIX) and prednisolone (XX). In many steroids there are both primary, secondary and tertiary alcohol functions which may all be transformed into esters. The selectivity, however, is reasonably good and the primary alcohol may be esterified by means of DCC as a coupling agent or by the direct use of the fatty acid chloride.



Scheme 5.

The thio derivatives of this invention may be prepared by methods analogous to those shown by the above reaction schemes.

The preparation of specific lipophilic antiinflammatory drug derivatives in accordance with the invention is illustrated by the Examples which follow, and in which Examples 6 and 7 illustrate the preparation of intermediate compounds.

EXAMPLE 1

2-Hydroxy-benzoic acid-(cis-9'-octadecenyl)ester

- 5 To a suspension of sodium hydride (60%) (0.21 g,
5.25 x 10⁻³ mol) in 40 ml anhydrous N,N-dimethylformamide
was added 2-hydroxy-benzoic acid (salicylic acid)
(0.726 g, 5.25 x 10⁻³ mol) and the mixture was stirred at
80°C under N₂ for 1 hour. Cis-9-octadecenol-mesylate
10 (1.82 g, 5.25 x 10⁻³ mol) was added and stirring was
continued for 22 hours. The cooled reaction mixture was
concentrated and the residue was dissolved in 100 ml
chloroform. The organic phase was washed with water,
diluted sodium bicarbonate and brine. The dried phase
15 was evaporated to dryness, and the crude product was
purified on a column of silica gel with 5% ether in
hexane as the eluent system. Homogenous fraction were
evaporated to give 1.3 g (64%) of the title compound.
¹H NMR(CDCl₃, 300MHz) δ: 10.85(1H, s, OH), 7.85(1H, d,
20 ArH), 7.45(1H, t, ArH), 6.95(1H, d, ArH), 6.85(1H, t,
ArH), 5.35(2H, m, CH=CH), 4.32(2H, t, CH₂-OCO),
1.95(4H, m, CH₂-C=), 1.75(2H, m, CH₂-C-O), 1.25(22H, m,
CH₂), 0.85(3H, t, CH₃).

25

30

35

EXAMPLE 2

2-(cis-9'-octadecenoxy)-ethyl-benzoate

5 To a suspension of sodium hydride (60%) (0.206g, 5.15×10^{-3} mol) in 40 ml anhydrous N,N-dimethylformamide was added 2-hydroxy-ethyl-benzoate (0.86g, 5.15×10^{-3} mol) and the mixture was stirred at 80°C under N₂ for 1 hour.

10 Cis-9-octadecenol-mesylate (1.78g, 5.15×10^{-3} mol) was added and stirring was continued for 40 hours. The cooled reaction mixture was evaporated at high vacuum, and the residue was treated with chloroform and water. The dried organic phase was concentrated and the crude

15 product was purified on a column of silica gel eluted with 5% ether in hexane. Homogenous fractions were collected to give 1.11g (52%) of the title compound.

¹H NMR (CDCl₃, 300 MHz) δ : 7.75(1H, d, ArH), 7.42(1H, t, ArH), 6.95(2H, m, ArH), 5.35(2H, m, CH=CH), 4.35(2H, q, CH₂-OCO), 4.0(2H, t, CH₂-OAr), 1.95(4H, m, CH₂-C=), 1.8(2H, m, CH₂-C-OAr), 1.48(2H, m, -CH₂-), 1.35(3H, t, CH₃-C-OCO), 1.25(20H, m, CH₂), 0.85(3H, t, CH₃).

20 ¹³C NMR (CDCl₃, 75 MHz) δ : 165.5(COO), 158.43(Ar C-2), 132.98(Ar C-4), 131.38(Ar C-6), 129.83 and 129.69(C=C), 120.81(Ar C-1), 119.83(Ar C-5), 112.97(Ar C-3), 68.75(CH₂-OAr), 60.58(CH₂-OCO), 31.82, 29.68, 29.40, 29.23, 29.16, 27.12, 25.92, 22.59(CH₂), 14.22(CH₃-C-OCO), 14.00(CH₃).

30

35

EXAMPLE 3

2-(cis-9'-octadecenoxo)-benzoic acid

- 5 To a suspension of 2-(cis-9'-octadecenoxo)-ethyl-benzoate (1.11g, 2.66×10^{-3} mol) in 25 ml ethanol and 50 ml water was added lithium hydroxide (2.0g) and the reaction mixture was stirred at 90°C for 6 hours. The ethanol was distilled off and 100 ml chloroform was added. pH
- 10 was adjusted to 7 with careful addition of 5N HCl, and the organic phase was washed with water. After removal of the solvent, the product was purified on a column of silica gel with 2% methanol in chloroform as the eluent system. 1.0g (96%) of the title compound was obtained.
- 15 ^1H NMR (CDCl_3 , 300 MHz) δ : 8.15(1H, d, ArH), 7.53(1H, t, ArH), 7.10(1H, t, ArH), 7.03(1H, d, ArH), 5.35(2H, m, CH=CH), 4.25(2H, t, $\text{CH}_2\text{-OAr}$), 1.95(4H, m, $\text{CH}_2\text{-C=}$), 1.92(2H, m, $\text{CH}_2\text{-C-OAr}$), 1.55-1.15(22H, m, $-\text{CH}_2-$), 0.85(3H, t, CH_3).
- 20 ^{13}C NMR (CDCl_3 , 75 MHz) δ : 165.44(COO), 157.48(Ar C-2), 134.81(Ar C-4), 133.42(Ar C-6), 129.80 and 129.51(C=C), 121.79(Ar C-5), 117.50(Ar C-1), 112.47(Ar C-3), 70.05($\text{CH}_2\text{-OAr}$), 31.73, 29.59, 29.52, 29.36, 29.15, 29.02, 28.98, 28.75, 27.04, 26.99, 25.58, 22.50(CH_2),
- 25 13.93(CH_3).

EXAMPLE 4

1-(p-Chlorobenzoyl)-5-methoxy-2-methylindole-3-acetic
acid-(cis-9'-octadecenyl)amide

5

- To a solution of 1-(p-chlorobenzoyl)-5-methoxy-2-methylindole-3-acetic acid (Indomethacin (0.56g, 1.56×10^{-3} mol) and TBTU (0.51g, 1.56×10^{-3} mol) in 6 ml anhydrous N,N-dimethylformamide was added
- 10 N,N-diisopropyl ethylamine (0.53 ml, 3.12×10^{-3} mol) and the reaction mixture was stirred under N₂ at room temperature for 30 minutes. A solution of cis-9-octadecenyl-amin (0.42g, 1.56×10^{-3} mol) in 6 ml anhydrous N,N-dimethylformamide was added and stirring
- 15 was continued for 3 hours. The solvent was evaporated at high vacuum and the residue was partitioned between chloroform and water. The dried organic phase was concentrated and the product purified on a column of silica gel with 2% methanol in chloroform as the eluent
- 20 system. Homogenous fractions were evaporated to give 1.05g of the title compound containing some DMF. The product was dissolved in ether, washed with water and the organic phase was dried and evaporated to give 0.88g (92%) of the title compound.
- 25 ¹H NMR (CDCl₃, 300 MHz) δ : 7.68(2H, d, ArH), 7.48(2H, d, ArH), 6.85(2H, m, ArH), 6.70(1H, dd, ArH), 5.60(1H, br. t, NHCO), 5.35(2H, m, CH=CH), 3.85(3H, s, CH₃O-Ar), 3.65(2H, s, Ar-CH₂-CO), 3.18(2H, q, CH₂-NH-), 2.40(3H, s, CH₃-Ar), 1.95(4H, m, CH₂-C=), 1.1-1.4(24H, m, CH₂),
- 30 0.85(3H, t, CH₃).

EXAMPLE 5

S(+)-2-(6-methoxy-2-naphthyl)propionic acid-(cis-9'-octadecenyl)-amide (naproxen oleyl amide)

5

- To a solution of naproxen (1.65g, 7.15×10^{-3} mol) and TBTU (2.30g, 7.15×10^{-3} mol) in 20 ml anhydrous N,N-dimethylformamide was added N,N-diisopropylethylamine (2.45 ml, 14.3×10^{-3} mol) and the reaction mixture was stirred under N₂ at room temperature for 30 minutes. A solution of 1-amino-cis-9-octadecen (1.91g, 7.15×10^{-3} mol) in 25 ml anhydrous N,N-dimethylformamide was added and stirring was continued for 3 hours. The solvent was evaporated at high vacuum and the residue was partitioned between chloroform and water. The dried organic phase was concentrated and the product purified on a column of silica gel with 3% methanol in chloroform as the eluent system. Homogenous fractions were evaporated to give 2.77g (81%) of the title compound.
- ¹H NMR (CDCl₃, 300 MHz) δ : 7.68(3H, m, ArH), 7.35(1H, d, ArH), 7.12(2H, m, ArH), 5.35(3H, m, CH=CH and NHCO), 3.92(3H, s, CH₃-OAr), 3.65(1H, q, CH), 3.15(2H, dt, CH₂-NHCO), 1.95(4H, m, CH₂-C=), 1.6(3H, d, CH₃), 1.25(24H, m, CH₂), 0.85(3H, t, CH₃).
- ¹³C NMR (CDCl₃, 75 MHz) δ : 174.09(CONH), 157.58(Ar C-6), 136.59(Ar C-10), 133.59(Ar C-9), 129.82 and 129.67(C=C), 129.02(Ar C-8), 128.85(Ar C-2), 127.35(Ar C-1), 126.22(Ar C-3), 125.96(Ar C-4), 119.00(Ar C-7), 105.47(Ar C-5), 55.16(CH₃-OAr), 46.92(CH), 39.55(CH₂-NH), 31.80, 29.66, 29.62, 29.40, 29.29, 29.22, 29.08, 27.10, 26.67, 22.58(CH₂), 18.43(CH₃-CH), 14.03(CH₃-CH₂).
- 10
- 15
- 20
- 25
- 30

EXAMPLE 6

S(+)-2-(6-hydroxy-2-naphthyl)propionic acid

- 5 To a well stirred suspension of sodium hydride (60%)
(12.9g, 0.336 mol) in 150 ml anhydrous
N,N-dimethylformamide was added dropwise a solution of
ethanethiol (24.3 ml, 0.328 mol) in 300 ml
10 N,N-dimethylformamide. A solution of naproxen (15g,
0.065 mol) in 150 ml N,N-dimethylformamide was slowly
added, and the reaction mixture was heated at 150°C for 3
hours. The clear solution was cooled and pH adjusted
(2-3) with 3.5N HCl. The solvents were evaporated at
15 high vacuum, and the residue was treated with a mixture
of 150 ml ether and 90 ml water. The solid deposit was
filtered off, and the filtrate concentrated. The residue
was treated with a mixture of 90 ml chloroform and 90 ml
water, and kept in the refrigerator for 24 hours.
20 The white deposit was filtered off, washed and dried to
yield 10.1g (72%) of the title compound.
¹H NMR (DMSO-d₆, 300 MHz) δ: 12.25(1H, s, COOH),
9.65(1H, s, Ar-OH), 7.75(1H, d, ArH), 7.65(2H, m, ArH),
7.31(1H, d, ArH), 7.05(2H, m, ArH), 3.75(1H, q, CH),
25 1.45(3H, d, CH₃).

EXAMPLE 7

S(+)-2-(6-hydroxy-2-naphthyl)propionic acid-ethyl ester

- 5 To a solution of S(+)-2-(6-hydroxy-2-naphthyl)propionic
acid (5.0g, 23×10^{-3} mol) in 1200 ml anhydrous ethanol
was added p-toluene-sulfonic acid (0.2g) and the reaction
mixture was heated at reflux for 24 hours. The cooled
mixture was stirred with a portion of solid NaHCO_3 . The
10 solution was filtered and the solvent was evaporated off.
The residue was dissolved in chloroform and washed with
water. The organic phase was concentrated, and the
crude product eluted through a column of silica gel with
2% methanol in chloroform. Homogenous fractions gave
15 4.8g (80%) of the title compound.
- ^1H NMR (CDCl_3 , 300 MHz) δ : 7.65(3H, m, ArH), 7.35(1H,
dd, ArH), 7.05(2H, m, ArH), 5.25(1H, br. s, Ar-OH),
4.15(2H, q, $\text{CH}_2\text{-OCO}$), 3.82(1H, q, CH), 1.58(3H, d, CH_3),
1.23(3H, t, $\text{CH}_3\text{-C-O}$).

EXAMPLE 8

S(+)-2-(6-[cis-9'-octadecenoxyl]-2-naphthyl)-propionic acid-ethyl ester

5

To a suspension of sodium hydride (60%) (0.47g, 11.8×10^{-3} mol) in 350 ml anhydrous N,N-dimethylformamide was added S(+)-2-(6-hydroxy-2-naphthyl)propionic acid-ethyl ester, and the reaction mixture was stirred under N₂ at room temperature for 2 hours. A solution of cis-9-octadecenol-mesylate (3.91g, 10.7×10^{-3} mol) in 5 ml N,N-dimethylformamide was added, and the stirring continued for 48 hours. The solvent was evaporated at high vacuum, and the residue was treated with chloroform and water. The dried organic phase was concentrated and the crude product was purified on a column of silica gel eluted with chloroform. Homogenous fractions gave 2.93g (56%) of the title compound.

¹H NMR (CDCl₃, 300 MHz) δ : 7.65(3H, m, ArH), 7.40(1H, d, ArH), 7.10(2H, m, ArH), 5.35(2H, m, CH=CH), 4.12(2H, q, CH₂-OCO), 4.05(2H, t, CH₂OAr), 3.82(1H, q, CH), 1.95(4H, m, CH₂-C=), 1.85(2H, m, CH₂-C-OAr), 1.55(3H, d, CH₃-CH), 1.45-1.20(22H, m, CH₂), 1.20(3H, t, CH₃-C-O), 0.85(3H, t, CH₃-CH₂).

¹³C NMR (CDCl₃, 75 MHz) δ : 174.67(COO), 157.08(Ar C-6), 135.65(Ar C-10), 133.67(Ar C-9), 129.94 and 129.79(C=C), 129.14(Ar C-8), 128.80(Ar C-2), 127.01(Ar C-1), 126.11(Ar C-3), 125.84(Ar C-4), 119.23(Ar C-7), 106.32(Ar C-5), 67.98(CH₂-OAr), 60.70(CH₂-OCO), 45.45(CH), 31.89, 29.74, 29.50, 29.46, 29.38, 29.31, 29.22, 27.18, 26.08, 22.67(CH₂), 18.59(CH₃-CH), 14.10(CH₃-CH₂- and CH₃-C-O).

EXAMPLE 9

S(+)-2-(6-[cis-9'-octadecenoxy]-2-naphthyl)-propionic
acid (naproxen oleyl ether)

5

A solution of S(+)-2-(6-[cis-9'-octadecenoxy]-2-naphthyl)-propionic acid-ethyl ester (3.79g, 7.67×10^{-3} mol) in 115 ml tetrahydrofuran and 25 ml 1M NaOH was stirred at room temperature for 10 days. 17 ml 1M HCl was added and the solvents were evaporated off.

10

The residue was taken up in chloroform and water, and the pH was adjusted to 1 with 1M HCl. The organic phase was washed with water, dried (MgSO_4) and concentrated to give 3.25g (94%) of the title compound.

15

^1H NMR (CDCl_3 , 300 MHz) δ : 7.61(3H, m, ArH), 7.35(1H, d, ArH), 7.10(2H, m, ArH), 5.35(2H, m, CH=CH), 4.03(2H, t, $\text{CH}_2\text{-OAr}$), 3.80(1H, q, CH), 1.95(4H, m, $\text{CH}_2\text{-C=}$), 1.82(2H, m, $\text{CH}_2\text{-C-OAr}$), 1.52(3H, d, $\text{CH}_3\text{-CH}$), 1.55-1.20(22H, m, CH_2), 0.85(3H, t, $\text{CH}_3\text{-CH}_2$).

20

^{13}C NMR (CDCl_3 , 75 MHz) δ : 180.96(COOH), 157.07(Ar C-6), 135.19(Ar C-10), 133.72(Ar C-9), 129.95 and 129.80(C=C), 129.17(Ar C-8), 128.76(Ar C-2), 127.01(Ar C-1), 126.17(Ar C-3), 126.02(Ar C-4), 119.18(Ar C-7), 106.30(Ar C-5), 67.98($\text{CH}_2\text{-OAr}$), 45.57(CH), 31.90, 29.76, 29.49, 29.43, 29.32, 29.25, 27.20, 26.11, 22.68(CH_2), 18.18($\text{CH}_3\text{-CH}$), 14.11($\text{CH}_3\text{-CH}_2$).

25

EXAMPLE 10

4-O-(trans-9'-octadecenoyl)-2-methyl-N(2-pyridyl)-2H-1,2-benzothiazine-3-carboxamide-1,1-dioxide

5

To a solution of 4-hydroxy-2-methyl-N[2-pyridyl]-2H,1,2-benzothiazine-3-carboxamide-1,1-dioxide (piroxicam) (2.5g, 7.54×10^{-3} mol) in 25 ml anhydrous N,N-dimethylformamide was added 2 ml of a solution of trans-9-octadecenoylchloride (2.2g, 7.53×10^{-3} mol) in 20 ml dichloromethane, and the reaction mixture was stirred under N₂ at room temperature. The remaining acid chloride solution was added in 2 ml portions at 2 hours intervals. After a total of 80 hours reaction time the solvents were evaporated off at high vacuum. The residue was dissolved in 200 ml ether and washed with water and a small amount of NaHCO₃(aq.). The dried (MgSO₄) organic phase was concentrated, and the crude product was purified on a column of silica gel eluted with ethylacetate/hexane (40:60). Homogenous fractions were collected and evaporated to give 3.56g of a solid material that was refluxed in pentane/ether, and the cooled mixture was kept at 4°C overnight. The solid material was filtered off, washed with pentane and dried to give 3.5g (78%) of the title compound.

¹H NMR (DMSO-d₆, 300 MHz) δ: 10.9(1H, s, NH) 8.38(1H, d, ArH), 8.08(1H, d, ArH), 7.7-8.0(5H, m, ArH), 7.20(1H, br. t, ArH), 5.35(2H, m, CH-CH), 3.1(3H, s, N-CH₃), 2.61(2H, t, CH₂-COO), 1.95(4H, m, CH₂-C=), 1.45(2H, m, CH₂-C-COO), 0.95-1.4(20H, m, CH₂), 0.85(3H, t, CH₃).

¹³C NMR (DMSO-d₆, 75 MHz) δ: 170.86(COO), 158.44(CONH), 150.93(C-2 pyr.), 148.10(C-6, pyr.), 138.33(C-4 pyr.), 135.30(C-4), 132.84(C-9), 131.67(C-6), 130.84(C-7), 130.03 and 130.01(C=C), 128.72(C-3), 128.49(C-10), 124.45(C-8), 122.21(C-5), 120.51(C-5 pyr.), 114.35(C-3 pyr.), 34.65(N-CH₃), 33.28, 31.97, 31.28, 29.02, 28.93, 28.84, 28.71, 28.51, 28.40, 28.25, 24.17, 22.10(CH₂), 13.91(CH₃).

EXAMPLE 11

[2-(2,6-Dichlorophenyl)amino]benzeneacetic acid)-(cis-9'-octadecenyl)-ester

5

To a solution of (2-[2,6-dichlorophenyl]-amino]benzeneacetic acid sodium salt) (diclofenac) (0.48g, 1.6×10^{-3} mol) in 15 ml dichloromethane and 3 ml N,N-dimethylformamide was added acetic acid (0.09 ml, 1.6×10^{-3} mol), cis-9-octadecen-1-ol (0.42g, 1.6×10^{-3} mol), 4-dimethyl-aminopyridine (DMAP) (50 mg) and DCC (0.34g, 1.7×10^{-3} mol), and the reaction mixture was stirred at 0°C for 6 hours and at a room temperature for 48 hours. A white deposit was filtered off and washed with dichloromethane. The organic phase was washed with water, dried (MgSO₄), concentrated and purified on a column of silica gel eluted with ethylacetate/hexane (40:60). Homogenous fractions gave 0.45g (53%) of the title compound as a colourless liquid.

20 ¹H NMR (CDCl₃, 300 MHz) δ: 7.35(2H, m, ArH), 7.25(1H, ArH), 7.15(1H, m, ArH), 6.95(2H, m, ArH), 6.58(1H, m, ArH), 5.35(2H, m, CH=CH), 4.15(2H, t, CH₂-O), 3.82(1H, s, Ar-CH₂-COO), 2.0(4H, m, CH₂-C=), 1.65(2H, m, CH₂-C-O), 1.25(22H, m, CH₂), 0.95(3H, t, CH₃).

EXAMPLE 12

4-O-(cis-11'-eicosenoyl)-2-methyl-N(2-pyridyl)-2H-1,2-benzothiazine-3-carboxamide-1,1-dioxide

5

- To a solution of 4-hydroxy-2-methyl-N[2-pyridyl]-2H-1,2-benzothiazine-3-carboxamide-1,1-dioxide (piroxicam) (0.3g, 0.990×10^{-3} mol) in 3 ml anhydrous N,N-dimethylformamide was added 1.5 ml of a solution of
- 10 cis-11-eicosenoylchloride (0.29g, 0.90×10^{-3} mol) in 2.5 ml dichloromethane, and the reaction mixture was stirred under N₂ at room temperature. The remaining acid chloride solution was added after 2 hours. After a total of 80 hours reaction time the solvents were evaporated
- 15 off at high vacuum. The residue was dissolved in 40 ml ether and washed with water and a small amount of NaHCO₃(aq.). The dried (MgSO₄) organic phase was concentrated, and the crude product was purified on a column of silica gel eluted with ethylacetate/hexane
- 20 (40:60). Homogenous fractions were collected and evaporated to give 0.42g (75%) of the title compound.
- ¹H NMR (DMSO-d₆, 300 MHz) δ : 10.9(1H, s, NH), 8.38(1H, d, ArH), 8.08(1H, d, ArH), 7.7-8.0(5H, m, ArH), 7.20(1H, br. t, ArH), 5.35(2H, m, CH=CH), 3.1(3H, s, N-CH₃),
- 25 2.61(2H, t, CH₂-COO), 1.95(4H, m, CH₂-C=), 1.45(2H, m, CH₂-C-COO), 0.95-1.4(24H, m, CH₂), 0.85(3H, t, CH₃).
- ¹³C NMR (DMSO-d₆, 75 MHz) δ : 170.84(COO), 158.43(CONH), 150.92(C-2 pyr.), 148.19(C-6 pyr.), 138.31(C-4 pyr.), 135.30(C-4), 131.65(C-6), 130.82(C-7), 129.58(C=C),
- 30 128.69(C-3), 128.46(C-10), 1214.43(C-8), 122.19(C-5), 120.47(C-5 pyr.), 114.33(C-3 pyr.), 34.65(N-CH₃), 33.29, 31.28, 29.11, 28.84, 28.70, 28.60, 28.27, 26.58, 24.17, 22.09(CH₂), 13.89(CH₃).

EXAMPLE 13

S(+)-2-(6-methoxy-2-naphthyl)propionic acid-cis-9'-octadecenyl-ester

5

To a solution of S(+)-2-(6-methoxy-2-naphthyl)propionic acid (naproxen) (0.15g, 0.65 mmol) in 10 ml dichloromethane was added cis-9-octadecenol (0.18g, 0.67 mmol), DCC (0.13g, 0.67 mmol), 4-dimethylaminopyridine (DMAP) (20 mg) and the reaction mixture was stirred under N₂ at room temperature for 3 hours. The white deposit formed was filtered off and washed with dichloromethane.

The solvent was evaporated off and the product was purified on a column of silica gel with dichloromethane as the eluent. Homogenous fractions gave 0.25g (80%) of the title compound.

¹H NMR (CDCl₃, 300 MHz) δ: 7.7(3H, m, ArH), 7.42(1H, d, ArH), 7.08(2H, m, ArH), 5.35(2H, m, CH=CH), 4.07(2H, t, CH₂-OCO), 3.9(3H, s, CH₃-OAr), 3.87(1H, q, CH), 1.95(2H, m, CH₂-C=), 1.25(22H, m, CH₂), 0.85(3H, t, CH₃).

20

EXAMPLE 14

11 β ,17 α ,21-trihydroxypregna-1,4-diene-3,20-dione-21-elaidate

5

- To a solution of 11 β ,17 α ,21-trihydroxypregna-1,4-diene-3,20-dione (prednisolone) (6.0g, 15.9 mmol) in 200 ml anhydrous dioxane and 6.5 ml pyridine was added elaidic acid chloride (8.0g, 26.6 mmol) and the reaction mixture was stirred at 10°C for 3 hours. A small amount of methanol was added and the solvents were evaporated at high vacuum. The residue was partitioned between ether and water. The organic phase was washed with tartaric acid (aq), NaHCO₃ (aq) and water. The dried organic phase was concentrated and the product purified on a column of silica gel with heptane/ethylacetate/methanol (64:32:4) as the eluent system. Homogenous fractions were evaporated to give 9.18g (90%) of the title compound.
- ¹H NMR (CDCl₃, 300 MHz) δ : 7.25(1H, br, d, CH=), 6.25(1H, dd, CH=), 6.0(1H, br, s, CH=), 5.38(2H, m, CH=CH), 4.92(2H, q, CH₂), 2.41(2H, t, CH₂-CO), 1.95(4H, m, CH₂-CH=), 0.87(3H, t, CH₃), 2.8 - 0.95(42H, m).

20

EXAMPLE 15

9-fluoro-11 β ,17,21-trihydroxy-16 β -methylpregna-1,4-diene-3,20-dione-21-elaidate

5

To a suspension of 9-fluoro-11 β ,17,21-trihydroxy-16 β -methylpregna-1,4-diene-3,20-dione (betamethasone) (0.9g, 2.3 mmol) in 40 ml anhydrous dioxane and 1 ml pyridine was added elaidic acid chloride (1.13g, 3.03 mmol) and the reaction mixture was stirred at ambient temperature for 48 hours. A small amount of methanol was added and the solvents were evaporated at high vacuum. The residue was partitioned between ether and water. The organic phase was washed with tartaric acid (aq), NaHCO₃ (aq) and water. The dried organic phase was concentrated and the product purified on a column of silica gel with heptane/ethylacetate/methanol (64:32:4) as the eluent system. Impure fractions were repurified and homogenous fractions were evaporated to give 1.02g (65%) of the title compound.

10

15

20

¹H NMR (CDCl₃, 300 MHz) δ : 7.22(1H, d, CH=), 6.31(1H, dd, CH=), 6.10(1H, br, s, CH=), 5.38(2H, m, CH=CH), 4.92(2H, q, CH₂), 2.41(2H, t, CH₂-CO), 1.95(4H, m, CH₂-CH=), 1.18(3H, d, CH₃), 0.87(3H, t, CH₃), 2,8 - 0.95(40H, m).

The lipophilic derivatives of the present invention may be administered systemically in the treatment of conditions for which NSAIDs and other antiinflammatory drugs are conventionally prescribed, either enterally or parenterally.

When administered enterally, which is the preferred form, the compounds of the invention may be formulated e.g. as soft or hard gelatine capsules, tablets, granules, grains or powders, dragees, syrups, suspensions or solutions.

When administered parenterally, preparations of the compounds of the invention as injection or infusion solutions, suspensions or emulsions are suitable.

The pharmaceutical compositions of this invention may be prepared by the usual techniques. Thus, the preparations can contain inert or pharmacodynamically active additives. Tablets or granulates e.g. can contain, as conventional, binding agents, filler materials, carrier substances or diluents. Liquid preparations may be present, for example, in the form of a sterile solution.

Capsules can contain a filler material or thickening agent in addition to the active ingredient. Furthermore, flavour-improving additives as well as the substances usually used as preserving, stabilizing, moisture-retaining and emulsifying agents, salts for varying the osmotic pressure, buffers and other additives may also be present.

If desired the pharmaceutical preparation of the compounds of the invention can contain an antioxidant, e.g. tocopherol, N-methyl-tocopheramine, butylated hydroxyanisole, ascorbic acid or butylated hydroxytoluene.

The dosages in which the compounds according to this invention are administered will vary according to the nature of the disease being treated and the stage of its development, to the mode of use and the route of use, as well as to the requirements of the patient. In general a daily dosage for systemic therapy for an adult average patient will be about 0.1-100 mg/kg body weight/day, preferably 0.5-30 mg/kg/day.

10

The invention further relates to a method for the treatment of inflammatory, pain-causing and/or pyretic conditions, which comprises administering at least one compound of the invention to a human patient in need of such treatment.

15

The currently preferred antiinflammatory drug lipophilic derivatives of the present invention are those in which the parent drug is naproxen. In particular, we have found that naproxen oleyl ether, naproxen oleyl ester and naproxen oleyl amide show improved antiinflammatory effect in comparison with naproxen itself. In animal in vivo models, these derivatives showed an improved effect on the first phase of the inflammation with respect to a reduction in the granuloma fluid content. An even more astonishing effect was the reduction of granuloma tissue dry weight especially with naproxen oleyl amide. This means a reduction of tissue damage, such as is not achieved by therapy with known NSAIDs. The effect was of an order which one could only expect from a therapeutic dose of steroids. Also the cartilage breakdown, a severe side effect of NSAIDs, was reduced.

20

The combination of these findings, e.g. the directly improved effect seen by the reduction in granuloma dry weight, and the reduction in cartilage breakdown, significantly improves the therapeutic index of the naproxen derivatives. The animals treated with the derivatives were also considerably less aggressive than

25

30

35

those treated with the parent compounds. This strongly indicates that the derivatives are less potent with respect to induction of gastrointestinal side effects.

- 5 Without wishing to be bound by theory, it is speculated that the enhanced antiinflammatory effects could be due to their lipophilic nature causing enhanced uptake by cells or due to activities that are quite separate from those of naproxen. The fatty acid tails added to
10 naproxen may act as scavengers of reactive oxygen species (ROS) which could be antiinflammatory through a number of mechanisms. For example, tissue damage will be inhibited through protection of ROS sensitive protease inhibitors, there will be prevention of formation of
15 endogenous antigens generated by oxidative damage to proteins and protection of hyaluronic acid from depolymerisation will prevent generation of angiogenic factors.
- 20 On implanted cartilage, naproxen like other NSAIDs, tend to increase proteoglycan and collagen loss. By contrast, the present naproxen derivatives show no tendency to increase proteoglycan or collagen loss from cartilage. Since inhibition of cyclooxygenase is thought to be
25 responsible for the detrimental effects of NSAIDs on cartilage, and since the naproxen derivatives share with naproxen the ability to inhibit this enzyme, it is suggested that their increased size and lipophilic nature results in exclusion from cartilage matrix.
- 30 Experiments which illustrate the enhanced antiinflammatory effects of these naproxen derivatives will now be described in detail.

35 Biological Effects

The in vivo model of granuloma induced cartilage degradation which was employed involves the implantation

- of sterile cotton-wrapped rat femoral head cartilage subcutaneously on the backs of mice. The cotton provokes a granulomatous response with a demonstrable T cell involvement which leads to loss of matrix compounds from the implanted cartilage. As a means of testing potential antiarthritic agents, this model has several distinct advantages. It involves chronic erosive disease with quantitative biochemical end points for determining the loss of cartilage matrix.
- 5
- 10 Antiinflammatory activity can be judged from the wet and dry mass of the cotton granuloma, chondroprotective activity can be determined from glycosaminoglycan and hydroxyproline content (indicative of proteoglycan and collagen respectively) of the implanted cartilage.
- 15 The granuloma is discrete and can be removed for the appraisal of various mediators or enzymes as required.

Groups of female (n = 10) TO mice (21 ± 4 g) received a subcutaneous implant of cotton-wrapped rat femoral head cartilage. At 2 weeks the implants were removed. The cotton provoked a granulomatous response with concomitant release of proteoglycan from the implanted cartilage. The effects of daily oral administration of equimolar amounts of naproxen (30 mg/kg) and naproxen derivatives (60 mg/kg) on granuloma development and cartilage proteoglycan content were assessed. The compounds were formulated as liposomes with empty liposomes acting as vehicle control.

20

25

- 30 A 15 mg/ml liposomal formulation is prepared by the 1:1 (w/w) mixing of the specific lipid derivatives (in DMSO) and lecithin (in ethanol) in a glycerol/sterile water buffer with subsequent dialysis to remove the solvents. A 7.5 mg/ml liposomal formulation of the underivatised NSAID compounds is prepared by the addition of the specific compound to empty liposomes in glycerol/sterile water.
- 35

Results were analysed with INSTAT using Mann-Whitney and with p values corrected for ties. Values of $p < 0.05$ were taken as significant.

5 Compounds tested

Naproxen (VI), naproxen oleyl ether (XII), naproxen oleyl ester (VII) and naproxen oleyl amide (VIII), $R' = \text{cis} - \text{CH}_2(\text{CH}_2)_7\text{CH}=\text{CH}(\text{CH}_2)_7\text{CH}_3$ in compounds XII, VII and VIII.

10 Description of the Figures

Figure 1 shows the fluid mass of the granulomas. Mean fluid content of granulomas from liposome-treated control animals was 62.69 mg. A reduction was seen in all treatment groups (naproxen 12%, naproxen oleyl ether 9%, naproxen oleyl ester 14% and naproxen oleyl amide 22%). The result with naproxen oleyl amide was especially significant.

Figure 2 shows the granuloma tissue dry weight.

20 Tissue dry weight of granulomas from liposome-treated control animals was 14.36 mg. Naproxen appeared to have no effect on tissue dry mass. Reductions were seen with the remaining treatment groups (naproxen oleyl ether 16%, naproxen oleyl ester 12% and naproxen oleyl amide 38%).
25 Again, the reduction observed with the naproxen oleyl amide was greatest.

Figure 3 shows the glycosaminoglycan content of cotton- . wrapped cartilages that had been implanted subcutaneously
30 into mice for 2 weeks. Non-implanted control cartilages had a mean glycosaminoglycan content of 1168 mg. Implantation into liposome-treated control animals for 2 weeks resulted in 60% loss of glycosaminoglycan. With the exception of implants from naproxen oleyl ether
35 treated animals, implants from the remaining treatment groups tended to have less glycosaminoglycan than those of the liposome-treated control group (naproxen 16%, naproxen oleyl ester 12% and naproxen oleyl amide 11%).

Figure 4 shows the hydroxyproline content of cotton-wrapped cartilages that had been implanted subcutaneously into mice for 2 weeks. Non-implanted control cartilages had a mean hydroxyproline content of 329 mg. Implantation into liposome-treated control animals for 2 weeks resulted in 19% loss of hydroxyproline. With the exception of implants from naproxen oleyl ether treated animals, implants from the remaining treatment groups tended to have less hydroxyproline than those of the liposome-treated control group (naproxen 12%, naproxen oleyl ester 8% and naproxen oleyl amide 3%).

The results obtained with naproxen in this model parallels what is known from similar studies with naproxen and other NSAIDs reported in the literature.

Fluid content of the granulomas was reduced when drug treatments were compared with controls, and was found significant in the case of the naproxen oleyl amide. Tissue dry mass appeared unaffected by naproxen, whereas the lipophilic derivatives appeared to cause a reduction which was again significant in the case of the naproxen oleyl amide. These remarkable findings strongly indicate that the naproxen derivatives reduce cartilage breakdown compared to naproxen itself. This was in fact confirmed, as naproxen, in comparison with the liposome control and the lipidic derivatives, appeared to increase proteoglycan loss from implanted cartilage. The same findings were mirrored for collagen breakdown, assessed as hydroxyproline content, although implants from naproxen-treated animals tended to have less collagen but there were no statistically significant differences between treatment groups.

Moreover, the naproxen-treated animals showed aggressive behaviour that resulted in the loss of 4 out of 10 implants. No implants were lost as a result of similar

behaviour in the liposome-treated group or those treated with the naproxen derivatives. This suggests that the naproxen derivatives were better tolerated than the parent NSAID.

5

The above experiments demonstrate a considerable improvement in the biological properties of naproxen by derivatising in accordance with the present invention.

10 Effect of prednisolone and betamethasone and their derivatives on rat peritoneal monocytes/macrophages

Male rats were injected intraperitoneally at time 0 with 4 ml of the test compounds at 10 mg/ml, or vehicle only.

15 At 6, 12, 25, 48 and 72 hours post treatment the peritoneal cavity was lavaged with 40 ml saline, the isolated cells were washed, counted and differentiated. The cells were then stimulated with opsonized zymosan, N-formyl-L-leucyl-L-phenylalanine (fMLP) or phorbol
20 12-myristate 13-acetate (PMA) and the activity of cells was measured by chemoluminescence production over one hour.

The effect of the derivatives are clear cut and
25 surprising. As seen from Figure 5, the effect of prednisolone on the activity of zymosan stimulated cells is only obvious as a slight reduction of the chemoluminescence at 6 hours. For the prednisolone-elaidate, the activity of the inflammatory cells are
30 reduced compared to control for up to 48 hours post treatment. The effect is clearly enhanced and prolonged compared to the effect of prednisolone itself.

Another set of experiments were performed to further
35 investigate the effect of selected prednisolone derivatives. In this experiment, the anti-inflammatory effect of 7 different fatty acid esters of prednisolone was compared.

Male rats were injected intraperitoneally with the test compound at time 0 at a dosage of 10 mg/ml or with vehicle only. At 48 hours post treatment the peritoneal cavity was lavaged, the isolated cells were washed, counted and differentiated. The cells were then stimulated with opsonized zymosan, N-formyl-L-Leucyl-L-phenylalanine (fMLP) or phorbol 12-myristate 13-acetate (PMA) and activity of the cells was measured by chemiluminescence production over one hour.

The effect of the derivatives are clearcut and surprising. As seen from fig. 6, the effect of prednisolone-elaidate, as an example of one of the especially preferred fatty acids, is the best, with the chemiluminescence only slightly affected for the other fatty acid derivatives. For the prednisolone-elaidate, the activity of the inflammatory cells are reduced compared to control and the rest of the fatty acid derivatives for up to 48 hours post treatment.

As shown in Figure 7, the number of cells in the peritoneal washing was substantially reduced, apparent up to 48 hours post treatment.

The differentiation of the cells showed that the major effect was on the number of macrophages in peritoneal lavage, as seen in Figure 8. For prednisolone, the effect was much less pronounced, and only seen at 6 and 12 hours. Similar effects, although not so clear cut, was seen in the comparison between betamethasone and betamethasone-elaidate, measured as the number of macrophages in the peritoneal lavage, Figure 9.

To investigate the direct anti-asthmatic effect of prednisolone-elaidic acid ester, the test compound was evaluated in the airway hyperresponsiveness model.

Effect of a prednisolone elaidic acid ester on the
endotoxin induced airway changes in rats.

In a model of acute inflammatory airway changes in rats,
5 the animals are exposed to aerosolized endotoxin (LPS).
Within 90 minutes this results in a diffuse neutrophilic
inflammation of bronchi and bronchioli, with a massive
increase of the number of neutrophils in the
bronchoalveolar fluid and an increase in airway
10 responsiveness, a key feature of asthma. 10 male F344
rats were exposed to 100 µg/ml LPS for 30 minutes in a
chamber. 12 and 4 hours prior to aerosol exposure the
animals were treated with either prednisolone or the
prednisolone derivative by instillation, 3 mg/kg. 90
15 minutes after the end of the aerosol exposure, the
animals were prepared for assessment of airway
responsiveness to 5-hydroxytryptamine and evaluation of
airway inflammation. 5HT was added intravenously every 5
minutes until minimum 50% increase in lung resistance is
20 observed, and the amount of 5 HT necessary to increase
the lung resistance with 50%, PC₅₀RL was calculated.
Neither prednisolone nor the derivative did influence the
number of inflammatory cells in the bronchoalveolar
fluid. The airway responsiveness was affected to a
25 surprisingly high extent by the prednisolone-elaidic acid
ester, as seen in fig. 10. This could be of great
importance in the treatment of asthma.

Cancer Drugs

30 Successful chemotherapeutic treatment of cancer is
limited by several main obstacles of which some may
completely or partly be overcome. Every change to a drug
that assures more specific action will directly benefit
35 the patient. A prime demand is that the tumour in
question is sensitive to the offered treatment. This may
be largely dependent on the class of therapeutics and
mechanism of action, and may be evaluated through

in vitro experiments on biopsies/isolated tumour cells prior to onset of the actual treatment. There are also known methods where a tumour can be sensitised towards several drugs.


5

Chemotherapeutic drugs are by their nature toxic to cells. As long as the malignant cells are more sensitive to the drug, there is a favourable situation.


10 If there are some prevalence for accumulation of the drug in tumour tissue/cells the therapeutic potential is further improved. To further improve the therapeutic index, organ targeting may be a vital factor. Very often a primary tumour, especially in an early stage, or as metastasis from another tumour type, is confined to
15 selected tissues like liver, spleen, lung, brain, etc. If the nature of the drug, its formulation, or mode of administration directs it to selected tissues, it may result in a very selective tumour eradication.

20 The preferred anti-cancer derivatives of the present invention have an improved therapeutic index, as will be demonstrated by the tests described below.

The parent cancer compound may be any compound which can
25 be categorized as having useful properties in the treatment of malignant tumours and which possesses one or more derivatisable groups selected from alcohol, ether, phenyl, amino, amido, thiol, carboxylic acid and carboxylic acid ester groups. Some examples of currently
30 available anti-cancer drugs which can be derivatised in accordance with the present invention include:




MEGESTROL




MEDROXYPROGESTERONE

52




5


HEXESTROL



TRILOSTANE




AMINOGLUTETHIMIDE




10

EPITIOSTANOL




CALUSTERONE




15

PODOPHYLLINIC ACID
2-ETHYLHYDRAZIDE




PIRARUBICIN

20




DOXORUBICIN




DAUNORUBICIN

53




TAXOL

5




MOPIDAMOL

10




MITOXANTRONE




LONIDAMINE

15




ETOPOSIDE




EFLORNITINE


20




DEFOSFAMIDE




TRIMETREXATE




METHOTREXATE




DENOPTERIN




THIOGUANIN



THIAMIPRINE




MERCAPTOPURIN




DACARBAZINE

55




NIMUSTINE




CHLOROZOTOCIN

5




MELPHALAN




ESTRAMUSTIN

10



CYCLOPHOSPHAMIDE



CHLORAMBUCIL

15




20 TRIMETHYLOLMELAMINE

As illustrated above, a number of known anti-cancer drugs
 25 contain more than one derivatisable group of the above-
 defined kinds. In these cases, one or more of these
 functional groups can be replaced by a lipophilic group
 in accordance with the present invention, and where there

are two or more lipophilic groups these may be the same or different lipophilic groups.

The lipophilic anti-cancer derivatives of the present invention may be prepared by the general preparative methods already described.

For example, the reaction scheme below illustrates the formation of amides and carbamates from doxorubicin (XXI) and daunorubicin (XXII). The amino group of the mother drug(s) can selectively be derivatised to an amide or a carbamate by means of reaction with the acyl-thiazolidine-2-thione or alkyloxy-carbonyl-thiazolidine-2-thione reagents made from the fatty acid (RCOOH) or the fatty alcohol (R'OH).




Scheme 6.

Again, Scheme 7 below illustrates the derivatisation of the two anti-cancer alkylating agents chlorambucil XXIII) and melphalan (XXIV). The mono-functional chlorambucil can be esterified or transformed into an amide by a number of methods. The bi-functional melphalan however, can undergo a number of side reactions such as self-condensation or ring forming reactions. In the unprotected melphalan, the use of coupling reagents such as DCC or TBTU is of limited value, but the amine function can conveniently be transformed into the

57

corresponding amide by means of an acyl-thiazolidine-2-thione reagent.



5

Scheme 7.

The preparation of specific anti-cancer derivatives in accordance with the invention is illustrated by the Examples which follow. Examples 18 and 21 relate to the preparation of intermediates.

10

EXAMPLE 16

5 Chlorambucil-oleyl ester

To a solution of 4-[p-[bis(2-chlorethyl)amino]-phenyl]butyric acid (chlorambucil) (0.966g, 3.18 mmol) and oleyl-alcohol (0.1893g, 3.33 mmol) in 70 ml
10 dichloromethane was added DCC(0.72g, 3.5 mmol) and N,N-dimethyl-amino-pyridine (DMAP) (25 mg), and the reaction mixture was stirred at ambient temperature for 12 hours. The solid deposit was filtered off, and the residue dissolved in 50 ml CH₂Cl₂ and washed with water.
15 25 ml ether was added to the organic phase and the solid deposit was filtered off. The filtrate was evaporated and the residue was purified on a column of silica gel with CH₂Cl₂ as the eluent. Homogenous fractions were evaporated to give 1.0g (55%) of the title compound.
20 ¹H NMR (CDCl₃, 300 MHz) δ: 7.10(2H, d, ArH), 6.65(2H, d, ArH), 5.35(2H, m, CH=CH), 4.05(2H, t, -CH₂-OCO), 3.75-3.55(8H, m, Cl-CH₂CH₂-N), 2.55(2H, t, Ar-CH₂-), 2.32(2H, t, CH₂-COO), 1.95(4H, m, CH₂-C=), 1.90(2H, t, Ar-C-CH₂-), 1.60(2H, m, CH₂-C-COO), 1.25(22H, m, CH₂), 0.85(3H, t,
25 CH₃).

EXAMPLE 17

Elaidic acid melphalan amide

- 5 To a solution of L-3-[p-[bis(2-chlorethyl)amino]-phenyl]alanine (melphalan) (0.603g, 1.98 mmol) in 24 ml DMF, 4 ml water and 4 ml triethylamine was added a solution of 3-thiazolidine-2-thione-elaidylamide (0.617g, 1.61 mmol) in 12 ml DMF, and the reaction mixture was
- 10 stirred at room temperature in the dark for 1.5 hours. The solvents were evaporated at high vacuum and the residue was dissolved in 100 ml chloroform and washed with water at pH 5.5. The organic phases was washed with AgNO₃ (aq), water (pH 5.5) and saturated NaCl (aq).
- 15 The organic phase was evaporated to give 0.84g (75%) of the title compound.
- ¹H NMR (CDCl₃, 300 MHz) δ: 7.08(2H, d, ArH), 6.60(2H, d, ArH), 6.05(NH), 5.35(2H, m, CH=CH), 4.75(N-CH-COO), 3.75-3.55(8H, m, Cl-CH₂CH₂-N), 3.2-2.95(2H, m, Ar-CH₂),
- 20 2.15(2H, t, CH₂-CON), 1.95(4H, m, CH₂-C=), 1.55(2H, m, CH₂-C-CON), 1.25(20H, m, CH₂), 0.85(3H, t, CH₃).

EXAMPLE 18

3-Elaidoyl-1,3-thiazolidine-2-thione

- 5 A mixture of elaidic acid (2.0g, 7.1 mmol), DMAP (86 mg, 0.7 mmol), 1,3-thiazolidine-2-thione (1.0g, 8.4 mmol) and DCC (1.7g, 8.2 mmol) in dichloromethane (20 ml) was stirred under N₂ at 0°C for 1 hour and then at ambient temperature for another 5 hours. Additional DCC (41 mg, 10 0.2 mmol) was added, and the reaction was stirred at the same temperature for 2 hours. Work up followed by flash chromatography (SiO₂; carbon tetrachloride-chloroform 1:0, 1:1, 0:1) gave 2.56g (94%) of the title compound as a yellow waxy solid.
- 15 ¹H NMR (CDCl₃, 300 MHz) δ: 5.36(2H, m, CH=CH), 4.56(2H, d, CH₂-NCO-), 3.26(2H, t, CH₂-S), 3.24(2H, t, CH₂-CON), 1.94(4H, m, CH₂-C=), 1.65(2H, m, CH₂-C-CON), 1.24(20H, m, CH₂), 0.86(3H, t, CH₃).

EXAMPLE 19

Elaidic acid daunorubicin amide

- 5 Daunorubicin hydrochloride (250 mg, 0.44 mmol) and
3-elaidoyl-1,3-thiazolidine-2-thione (Example 18, 400 mg,
1.04 mmol) were partitioned between THF (20 ml) and brine
(20 ml 4M NaCl), buffered with sodium carbonate (0.12M
NaHCO₃, 0.8M Na₂CO₃). The mixture was vigorously
10 stirred in the dark under N₂ for 4 hours at ambient
temperature. The phases were separated, and the aqueous
phase extracted with ether (3 x 10 ml). The combined
organic phase was washed with aqueous sodium nitrate
(3 x 10 ml 2M). To remove the 1,3-thiazolidine-2-thione
15 present, pyridine (1.0 ml) was added and the ether phase
shaken vigorously with aqueous solution (2 x 3 ml) of
sodium nitrate (2M) containing silver nitrate (0.2M).
After each treatment the mixture was filtered through
Celite with ether (20 ml) used as rinse. The ether
20 phase was washed with aqueous sodium nitrate (5 ml 2M)
and brine (5 ml) and finally dried (MgSO₄). The crude
product was purified on a column of silica gel prepared
with pyridine (0.2% w/w) and with 0.2% pyridine and 0.6%
methanol in chloroform as the eluent to give 332 mg (95%)
25 of the title compound as a dark red powder.
- ¹H NMR (CDCl₃, 300 MHz) δ: 13.87(1H, s), 13.09(1H, s),
7.9(1H, d), 7.69(1H, t), 7.29(1H, d), 6.05(1H, d),
5.41(1H, s), 5.31(2H, m), 5.10(1H, s), 4.16(3H, m),
3.97(3H, s), 3.94(1H, m), 3.61(2H, m), 3.09(1H, d),
30 2.71(1H, d), 2.36(3H, s), 2.24(1H, d), 2.06(2H, m),
1.86(4H, m), 1.77(2H, m), 1.50(2H, m), 1.25(3H, d),
1.21(20H, m), 0.83(3H, t).

EXAMPLE 20

Elaidic acid doxorubicin amide

- 5 Doxorubicin hydrochloride (400 mg, 0.69 mmol) was treated as described above with 3-elaidoyl-1,3-thiazolidine-2-thione (Example 18, 400 mg, 1.04 mmol) in THF (35 ml) and buffered brine (35 ml) for 10 hours at ambient temperature. Additional amidation reagent (Example 18:
- 10 100 mg, 0.26 mmol) was necessary to complete the reaction. After 6 hours the phases were separated and the aqueous phase extracted with THF (2 x 15 ml). The combined organic phase was washed with brine and dried (MgSO₄). Purification by flash chromatography as
- 15 described in Example 19 afforded 440 mg (79%) of the title compound as dark red crystals (mp. 115-116°C).
- ¹H NMR (CDCl₃, 300 MHz) δ: 14.13(1H, s), 13.31(1H, s), 7.98(1H, d), 7.74(1H, t), 7.35(1H, d), 5.87(1H, d), 5.46(1H, d), 5.33(2H, m), 5.20(1H, s), 4.73(2H, s),
- 20 4.52(1H, s), 4.13(2H, m), 4.03(3H, s), 3.61(1H, m), 3.20(1H, d), 3.02(1H, br. s), 2.89(1H, d), 2.4-2.0(5H, m), 2.0-1.6(6H, m), 1.53(2H, m), 1.26(3H, d), 1.23(20H, m), 0.85(3H, t).

EXAMPLE 21

3-(cis-9-Octadecen-1-oxycarbonyl)-1,3-thiazolidine-2-thione

5

The compound was prepared essentially as described by Chen and Yang for the ethyl analogue. Oleyl alcohol (cis-9-octadecen-1-ol; 2.8g, 10.4 mmol) was added within 10 minutes to a stirred solution of 2-thioxo-3-thiazolidinecarbonyl chloride² (1.6g, 8.6 mmol) and TEA (1.3 ml, 9.3 mmol) in chloroform (dry, ethanol-free; 15 ml) kept at 0°C under N₂. The mixture was stirred at the same temperature for 80 minutes when ice-water (5 ml) was added. The pH of the aqueous phase was adjusted to 15 6 by dropwise addition of hydrochloric acid (0.5 ml 1M). Standard work up followed by flash chromatography (SiO₂: hexane-chloroform 1:1, 1:2, 1:3, 0:1) gave 1.78g (50%) of a yellow oil.

¹H NMR (CDCl₃, 300 MHz) δ : 5.32(2H, m, CH=CH), 4.50(2H, d, CH₂-NCO-), 4.25(2H, t, CH₂-OCO-), 3.28(2H, t, CH₂-S), 1.99(4H, m, CH₂-C=), 1.69(2H, quint., CH₂-C-OCON), 1.26(22H, m, CH₂), 0.86(3H, t, CH₃).

20

EXAMPLE 22

Daunorubicine oleyl carbamate, [N-(cis-9-octadecen-1-oxycarbonyl)daunorubicin]

5

Daunorubicin hydrochloride (250 mg, 0.44 mmol) was treated with 3-(cis-9-octadecen-1-oxycarbonyl)-1,3-thiazolidine-2-thione (Example 21, 550 mg, 1.33 mmol) for 27 hours as described for the amide analogue in Example

10 19. The crude product obtained after extractive work up with THF as described in Example 20 was dissolved in ether (40 ml). Removal of the 1,3-thiazolidine-2-thione present was achieved according to Claim 19. The crude product was purified on a column of silica gel prepared
15 with pyridine (0.2% w/w) and with 0.2% pyridine and 0-10% methanol in benzene as the eluent to give 321 mg (87%) of the title compound as a dark red powder.

¹H NMR (1% pyridine-d₆ in CDCl₃, 300 MHz) δ: 13.95(1H, s), 13.24(1H, s), 8.00(1H, d), 7.74(1H, t), 7.36(1H, d),
20 5.48(1H, d), 5.32(2H, m), 5.25(1H, s), 5.09(1H, d), 4.51(1H, br. s), 4.18(1H, m) 4.05(3H, s), 3.94(2H, t), 3.85(1H, m), 3.66(1H, s), 3.20(1H, d), 2.90(1H, d), 2.39(3H, s), 2.31(1H, d), 2.08(1H, dd), 1.97(4H, m), 1.9-1.6(3H, m), 1.51(2H, m), 1.28(3H, d), 1.23(22H, m),
25 0.84(3H, t).

EXAMPLE 23

Doxorubicin oleyl carbamate, [N-(cis-9-octadecen-1-oxycarbonyl)doxorubicin]

- 5
Doxorubicin hydrochloride (250 mg, 0.43 mmol) was treated with 3-(cis-9-octadecen-1-oxycarbonyl)-1,3-thiazolidine-2-thione (Example 21: 700 mg, 1.69 mmol) for 69 hours as described for the amide analogue in Example 19. THF was
10 removed at reduced pressure and the resulting suspension was partitioned between water (20 ml) and pyridine-chloroform 1:4 (25 ml). Still undissolved material was treated with pyridine (15 ml combined) added to the organic fraction. Volatiles were removed by evaporation
15 under reduced pressure. The resulting residue (1.1g) was sonicated with ethyl acetate containing an aqueous solution of silver nitrate (0.6 ml 1M) for 20 minutes at 20-30°C. The resulting suspension was filtered through Celite with ethyl acetate (20 ml combined) used as rinse.
20 The sonication cycle was repeated with further silver nitrate (0.4 ml 1M). The combined organic phase was washed with brine (5 ml) and dried (MgSO₄). The crude product (0.61g) obtained by evaporation of the ethyl acetate on a rotary evaporator was purified by flash
25 chromatography as described in Example 22 to give 208 mg (58%) of the title compound as a dark red glass.
- ¹H NMR (1% pyridine-d₆ in CDCl₃, 300 MHz) δ: 14.03(1H, s), 13.29(1H, s), 7.99(1H, d), 7.75(1H, t), 7.36(1H, d), 5.48(1H, d), 5.30(3H, m), 5.25(1H, s), 5.05(1H, d),
30 4.74(2H, s), 4.2-4.0(2H, m), 4.05(3H, s), 3.95(2H, t), 3.82(1H, m), 3.65(1H, s), 3.21(1H, d), 2.92(1H, d), 2.31(1H, d), 2.14(1H, dd), 1.97(4H, m), 1.9-1.7(3H, m), 1.52(2H, m), 1.28(3H, d), 1.24(22H, m), 0.85(3H, t).

EXAMPLE 24

Taxol-2'-elaidate

- 5 Taxol (25mg, 0.029 mmol) was initially dried by repeatedly dissolving in pyridine (3 x 1 ml) and evaporating under reduced pressure. It was then dissolved in pyridine (1 ml) to give a clear, colourless solution, to which was added N-(3-dimethylaminopropyl)-
10 N'-ethylcarbodiimide hydrochloride (9mg, 0.05 mmol), DMAP (2mg, 0.02 mmol), elaidic acid (10mg, 0.035 mmol) and anhydrous MgSO₄ (3mg) as a mixture. The reaction was then stirred for 48 h at room temperature under N₂. The pyridine was removed by evaporation under reduced
15 pressure and the residue dissolved in DCM (25 ml). The organic phase was then dried (MgSO₄), filtered and evaporated under reduced pressure to give a white solid, which was purified by flash chromatography (SiO₂; diethyl etherhexane 1:1 to 1:0 gradient elution) to yield the
20 title compound as a white solid (25mg, 77%).
- ¹H NMR (CDCl₃, 300 MHz) δ : 0.87(3H, t), 1.33(3H, s), 1.2-1.8(23H, m), 1.57(3H, t), 1.68(3H, s), 1.76(1H, s), 1.8-2.1(8H, m), 2.17(1H, m), 2.23(3H, s), 2.40(3H, m), 2.46(3H, s), 2.52(1H, d), 2.56(1H, m), 3.81(1H, d),
25 4.20(1H, d), 4.32(1H, d), 4.45(1H, m), 4.97(1H, d), 5.38(2H, dt), 5.50(1H, d), 5.68(1H, d), 5.94(1H, dd), 6.27(1H, t), 6.29(1H, s), 6.88(1H, d), 7.35-7.44(7H, m), 7.51-7.54(3H, m), 7.60-7.68(1H, m), 7.23(2H, d) and 8.13(2H, d).

Antitumour effect of melphalan-elaidic amide and
chlorambucil-oleylester in vivo using the murine
subcutaneous ADJ/PC5 plasmacytoma and its cisplatin-
5 resistant subline

Cytotoxicity of chlorambucil and chlorambucil-fatty acid
conjugates of different degree of unsaturation against
human lymphomas and normal human peripheral blood
10 lymphocytes has been described of A. Anel et al,
Biochemical Pharmacology, vol. 40, no. 6, pages
1193-1200, 1990. The toxicity of Chlorambucil-
arachidonic acid and Chlorambucil-docosahexaenoic acid
against lymphoma cells was equal or higher than the
15 individual toxic potential of either chlorambucil or the
free fatty acids. To the contrary, the fatty acid
derivatives of this invention, exemplified by oleic and
elaidic acid, are much less toxic than the mother drug
alone, as the following experiment shows.

20 Murine solid ADJ/PC6 plasmacytoma and its subline
selected for resistance to cisplatin and other alkylating
agents was implanted subcutaneously as 1mm³ tumour
fragments into BALB/C female mice weighing 20-25 grams.
25 Melphalan or melphalan-elaidic amide, or chlorambucil or
chlorambucil-oleylester, were administered by
intraperitoneal, single dose 20 days post subcutaneous
tumour implantation. Tumours were dissected on day 30,
and weights of control and treated groups were compared.
30 Activity was based on the measurement of drug toxicity,
LD₅₀ in mg/kg compared to anti-tumour effect measured as
ED₉₀, the dose in mg/kg required to reduce tumour mass by
90% compared to controls.

35 As can be seen from Table 1, a much greater dose was
necessary to achieve LD₅₀ in mg/kg for both melphalan-
elaidic amide compared to melphalan and for chlorambucil-
oleylester compared to chlorambucil. This means that

toxicity was reduced.

Table 1

	Mother compound LD ₅₀	Derivative LD ₅₀
Melphalan	23 mg/kg	180 mg/kg
Chlorambucil	57 mg/kg	>1600 mg/kg

5

ED₉₀ was achieved for melphalan-elaidic acid amide for both the sensitive and cisplatin resistant tumours, whilst no activity was shown for melphalan in the cisplatin resistant tumour, with ED₉₀ for melphalan-elaidic amide of 60 mg/kg.

10

Cellular accumulation of Doxorubicine and Doxorubicine-derivatives in cells, with or without multi-drug resistance

15

Tumour cells may become resistant to anti-cancer drugs after a prolonged chemotherapy. One form of drug resistance is the multi-drug-resistance (MDR) where the cells are cross-resistant to a variety of drugs such as vinca alkaloids, anthracyclines, actinomycin D and colchicine. The MDR phenotype has been correlated with the overexpression of a particular class of transmembrane glycoprotein called the P-glycoproteins. P-gp seems to function as an energy-dependent drug-efflux pump.

25 The P-glycoproteins can decrease the intracellular concentration of an anti-cancer drug below its active concentration by actively pumping the drug out of the cell. Verapamil, a calcium channel blocker can reverse MDR by increasing the intracellular concentration of

30 anti-tumour drug. Dihydropyridine and pyridine analogues, calmodulin inhibitors, synthetic isoprenoids, lysosomotropic agents, bisbenzylisoquinoline alkaloids, quinidine, quinacrine, lidocaine, phenoxazine, amiodarone and cyclosporin A are other examples of drugs which

35 alters the MDR when co-administered to cells or

co-administered in vivo. Although still at an experimental level, the use of resistance modulators in cancer therapy are becoming popular.

5 The co-administration of these highly bio-active compounds are not without problems themselves. Mild to severe, life threatening side effects, are observed, which prohibit the transfer of the very promising in vitro results to a clinical situation.

10 Most of these agents are cationic and lipophilic. Lipophilicity is a desirable characteristic for a resistance modulator. Liposome-encapsulated doxorubicin has also been tested for its effectiveness in overcoming multi-drug resistance. The intracellular concentration
15 of drug has been doubled using the liposomes of doxorubicin (Cancer chemotherapy and Pharmacology, 1991, 28: 259-265). Liposomes alone may also influence MDR (Increased accumulation of drugs in multi-drug resistant cells induced by liposomes, cancer research, 52,
20 3241-3245, 1992), (liposomes of cardiolipin, phosphatidylinositol, dioleoylphosphatidic acid).

Cells were exposed in suspension at 2×10^5 /ml to drugs at 20 μ M. Aliquots were taken at various times and
25 rinsed in ice cold PBS before being passed through the flow cytometre. Accumulation as a function of time of Doxorubicine and Doxorubicine-derivatives is seen in Figure 11. In cell line COR-L23/P (human large lung cell) and its resistant cell line COR-L23/R the
30 concentration of the Doxorubicine-derivatives (Doxorubicine-elaidic-amide and Doxorubicine-oleyl-carbamate) is approximately the same, whilst Doxorubicine concentration is much lower in the resistant cell line.

35 Cellular accumulation of Daunorubicine and Daunorubicine-derivatives in cells with or without multi-drug resistance

Cells were exposed to Daunorubicine, Daunorubicine elaidic amide and Daunorubicine oleic carbamate, as described above. Drug concentration was reduced to 10 μM . Fluorescence due to uptake in resistant and non-resistant cells are more or less equal in the two cell types for the derivatives, as seen in Figure 12, compared to the fluorescence due to uptake in the same cells for Daunorubicine itself (Figure 13).

10 Sensitisation of a cell to Doxorubicin by
15 co-administration of Doxorubicin-elaidic-amide

An MTT assay was used to determine the toxicity of compounds. The cells were exposed to the compounds for 6 days prior to assaying. The cell line used was H69/LX4, human small lung cell line, overexpressing PgP. The cell line is highly resistant to Doxorubicine-elaidic-amide alone, with an IC_{50} value of Doxorubicine-elaidic-amide $>50 \mu\text{M}$. But surprisingly, if Doxorubicine-elaidic-amide is administered at 5 μM in addition to Doxorubicine, the sensitivity of the cell line to Doxorubicine is enhanced from $\text{IC}_{50}=0.4 \mu\text{M}$ to $\text{IC}_{50}=0.08 \mu\text{M}$. Addition of 20 μM Doxorubicine-elaidic-amide enhances the sensitivity for Doxorubicine to $\text{IC}_{50}=0.04 \mu\text{M}$. The results given in Table 2 show that the derivative has the ability to interact with the resistance mechanism of the cells and potentiate and restore the effect of Doxorubicine itself down to the level of a sensitive cell line.

30
Table 2

Doxorubicine-elaidic amide	$\text{IC}_{50}\mu\text{M}$, Doxorubicine
0 μM	0.4 μM
5 μM	0.08 μM
20 μM	0.04 μM

As shown in Figures 11-13 above, fatty acid derivatives of the anthracyclines doxorubicin and daunorubicin have a modulating effect on the MDR mechanism of a dox resistant cell line. When co-administered with the mother drug to a resistant cell line, the sensitivity towards the mother drug is again at the same order of magnitude as in the sensitive line. This approach towards MDR modulators is favourable because the co-administered drug is just a derivative of the active compound, which when/if it is hydrolysed in vivo liberates the active drug and a non-toxic fatty acid residue.

Antimicrobial Agents

There is a vast variety of drugs falling within this therapeutic area.

The most important class of antimicrobial agents are probably the penicillins, but as drug resistance is becoming more serious, focus is directed towards alternative therapeutics for treating bacterial infections. Although treated with other drugs with other mechanisms of action, some of the same factors important to fighting bacterial infections may be valid for the treatment of diseases caused by mycobacterias and protozoa, for example factors such as cellular uptake, tissue distribution, circumvention of resistance mechanisms.


All the drugs used within this field have very good to moderate effect towards the target infectious species. Aside normal new drug development, much focus within this field has been directed towards the development of derivatives with better oral bioavailability, i.e. pure pro-drugs, and very little of this work has had any impact on drug resistance problems.

The clinical efficacy of an antibiotic is determined not


- only by its antibacterial activity but also by its pharmaceutical and pharmacokinetic properties. Pro-drugs have been used to increase the stability and solubility of antibiotics and further improve the oral absorption, tissue penetration and duration of the parent compounds. Enhanced serum levels after oral administration will lead to improved tissue concentrations of antibiotic.
- 10 Within the penicillin and other related β -lactam antibiotics field it is often stated that simple alkyl esters are too stable to be used as pro-drugs. The preferable pro-drugs can be double esters with one to three methylene linkers or methoxycarbonyl alkyl esters.
- 15 These side-chain modifications facilitate improved oral bioavailability, and the derivatives are sufficiently biolabile to release the active drug through hydrolysis catalysed by endogenous or microorganism-induced enzymes in the bloodstream. These penicillin pro-drugs have
- 20 little influence on the drug resistance situation. The most predominant and reasonably well characterised resistance mechanisms active towards penicillins and closely related analogues are the bacteria's acquired ability to produce the hydrolysing enzyme β -lactamase.
- 25 The enzyme can be found both intra- and extracellularly, meaning that the drugs can be broken down in the bloodstream even before they reach the actual bacteria. Other types of drug resistance may be due to a pure exclusion mechanism, whereby the drugs are prevented from
- 30 entering the bacteria or other microorganisms. The lipid derivatives of this invention are not so easily hydrolysed in the bloodstream, whereby better circulation of the drug-derivative in the bloodstream can be achieved. It is believed that the especially high
- 35 cellular transport and addition efficiency of the novel lipid derivatives overcomes the exclusion mechanisms, and the active drug is released in other compartments of the cells/bacteria where it can exert its action irrespective

of hydrolytic enzymes.


Some examples of antibiotics and other antibacterial agents which can be derivatised in accordance with this invention include the following compounds:




OXACILLIN




AMPICILLIN




AMOXICILLIN




CEPHALEXIN




CEPHALOTIN



CEPHALOSPORIN




DOXYCYCLIN




CHLORAMPHENICOL

74




p-AMINO-SALICYLIC ACID




ETHAMBUTOL

5




CIPROFLOXACIN




ENROFLOXACIN

10



DIFLOXACIN




DANOFLOXACIN

- 15 As illustrated above, antibacterial drugs may contain
 more than one derivatisable group, and in these cases one
 or more of these functional groups can be replaced by a
 lipophilic group in accordance with the present
 invention, and where there are two or more lipophilic
 20 groups these may be the same or different lipophilic
 groups.

The lipophilic antibacterial compounds of the present
 invention may be prepared by the same general methods as
 25 heretobove described. However, it is to be noted that
 the selective and efficient derivatisation of several
 penicillin derivatives may be complicated by various
 factors such as the presence of multiple reactive groups

(-OH₂-NH- and -NH₂) in the mother drugs, due to ring opening of the β -lactams, other rearrangements or breakdown of the compounds. The use of protective groups and various reagent systems may therefore facilitate selective derivatisation, as illustrated for ampicillin (XXV) in Scheme 8.


The primary amino group can selectively be transformed into fatty-acid ampicillin amide (XXVI) by means of the acyl-thiazolidine-2-thione. The same amino function can be protected as a Schiff base (XXVII) with benzaldehyde. The carboxylic acid is transformed into the Cesium salt and further reacted with the fatty bromide (RBr). Mild acidic hydrolysis reforms the amino function to give the ampicillin-fatty acid ester (XXVIII).



20 Scheme 8

Selective derivatisation of the tri-functional anti-tuberculosis agent para-amino-salicylic acid, PAS (XXIX) is shown in Scheme 9. PAS itself is unstable under a number of reaction conditions, and both self-condensation and the formation of di-adducts are possible. The carboxylic acid can be converted to its Caesium salt and

further transformed into the corresponding ester through the reaction with a mesylate. Fatty acid chlorides reacts primarily with the amino function to give the corresponding amide. Products resulting from reaction on the phenolic group can be removed through a slightly basic hydrolysis.



10 Scheme 9.

The preparation of specific antibacterial compounds in accordance with this invention is illustrated in the Examples which follow.

EXAMPLE 25

Para-amino-salicylic acid elaidylester

- 5 To a suspension of para-amino-salicylic acid (PAS)
(0.47g, 3.1 mmol) and Cesium carbonate (0.98g, 3.0 mmol)
in 50 ml anhydrous DMF was added elaidylmesylate (1.0g,
2.9 mmol) and the reaction mixture was stirred at ambient
temperature for 60 hours and at 35°C for 48 hours. The
10 reaction mixture was extracted with ether and water, and
the organic phase was washed with NaHCO₃(aq) and water.
The solvent was evaporated off and the residue was
purified on a column of silica gel with
heptane/CH₂Cl₂/AcOH/MeOH (85:15:2:2) as the solvent
15 system. Product containing fractions was repurified
(70:30:2:2) and homogenous fractions were evaporated to
give 0.6g (51%) of the title compound.
- ¹ H NMR (DMSO-d₆, 300 MHz) δ: 10.85(1H, s, -OH), 7.45(1H,
d, ArH), 6.15(1H, d, ArH), 5.95(1H, s, ArH), 5.35(2H, m,
20 CH=CH), 4.20(2H, t, CH₂-O), 3.5(2H, br. s, NH₂), 1.95(4H,
m, CH₂-C=), 1.65(2H, m, CH₂), 1.25(22H, m, CH₂), 0.85(3H,
t, CH₃).

EXAMPLE 26

4-(elauidamido)-salicylic acid

5

To a solution of PAS (2.6g, 17 mmol) in 100 ml anhydrous THF and 4 ml pyridine was added dropwise a solution of elaidic acid chloride (5.11 g, 17 mmol) in 20 ml THF at 0°C. The reaction mixture was stirred at ambient temperature for 24 hours. More elaidic acid chloride (1.0g) was added, and after 4 hours, a small amount methanol was added. The solvents were evaporated at high vacuum, and the residue was partitioned between ether and water. The organic phase was washed with tartaric acid (aq) and water. The solvent was evaporated off, and the residue was dissolved in 100 ml methanol to which was added 5 ml water. The solid deposit was filtered off and recrystallised from ethanol to give 3.9g crude material. 1.5g of this material was dissolved in 100 ml ethanol to which was added 11ml 1M NaOH(aq). The mixture was stirred at ambient temperature for 2 hours and acidified with 30 ml 0.5M tartaric acid(aq). The solid deposit was washed with water, dissolved in ether, and the organic phase washed with tartaric acid(aq). The solvent was evaporated off, and the residue was evaporated from anhydrous chloroform x 2, to give 1.3g(87%) of the title compound.

¹H NMR (DMSO-d₆, 300 MHz) δ: 11.5(1H, br. s, COOH), 10.15(1H, s, Ar-OH), 7.70(1H, d, ArH), 7.35(1H, s, ArH), 7.05(1H, dd, ArH), 5.35(2H, m, CH=CH), 2.31(2H, t, CH₂-CON), 1.95(4H, m, CH₂-C=), 1.55(2H, m, CH₂-C-CON), 1.25(20H, m, CH₂), 0.85(3H, t, CH₃).

EXAMPLE 27

Elaidic acid chloramphenicol ester.

- 5 To a solution of D-threo-2,2-dichloro-N-[β -hydroxy- α -(hydroxymethyl)]-p-nitrophenethylacetamide (0.20g, 0.62 mmol) in 10 ml anhydrous DMF and 2 ml pyridine was added a solution of elaidic acid chloride (0.19g, 0.62 mmol) in 3 ml DMF. The reaction mixture was stirred at ambient
10 temperature for 12 hours. The solvents were evaporated at high vacuum, and the residue was partitioned between ethyl acetate and water. The organic phase was concentrated and the crude product was purified on a column of silica gel with ethylacetate/hexane (1:1) as
15 the solvent system. Homogenous fractions were evaporated to give 0.14g(39%) of the title compound.

^1H NMR (DMSO- d_6 , 300 MHz) δ : 8.52(1H, m, NH), 8.17(2H, d, ArH), 7.12(2H, d, ArH), 6.42(1H, s, CHCl), 6.22(1H, d, OH), 5.35(2H, m, CH=CH), 5.00(1H, m, CH-O), 4.25 and
20 4.15(3H, m, CH-N and CH-OCO), 2.25(2H, t, CH_2COO), 1.95(4H, m, $\text{CH}_2\text{-C=}$), 1.25(20H, m, CH_2), 0.85(3H, t, CH_3).

EXAMPLE 28

Oxacillin-oleyl ester.

- 5 To a solution of (5-methyl-3-phenyl-4-isoxazoly)penicillin (oxacillin)sodium salt (1.0g, 2.4 mmol) in 76 ml water was added 13.1 ml 0.2M HCL(aq), and the mixture was evaporated. The residue was dissolved in 50 ml methanol and 5 ml water. A 20% solution of
10 Cs_2CO_3 (aq) was added until pH 7. The mixture was evaporated to dryness.

- To a solution of the residue in 50 ml DMF was added oleyl bromide(0.78g, 2.4 mmol) and the reaction mixture was stirred at ambient temperature for 72 hours. The solvent
15 was evaporated at high vacuum, and the residue was extracted with water and chloroform. The organic phase was concentrated, and the crude product was purified on a column of silica gel with ethylacetate/hexane (40:60) as the solvent system. Homogenous fractions were evaporated
20 to give 0.69g (45%) of the title compound.

- ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ : 9.3(1H, d, CO-NH), 7.7 and 7.5(5H, m, ArH), 5.6(2H, m, N-CHCH-S), 5.3(2H, m, CH=CH), 4.4(1H, s, CHCOO), 4.15(2H, t, COOCH₂), 2.55(3H, s, CH₃), 1.95(4H, m, CH₂-C=), 1.55(2H, m, CH₂-C-OOC), 1.52 and
25 1.42(6H, s, CH₃), 1.25(22H, m, CH₂), 0.85(3H, t, CH₃).

EXAMPLE 29

Elaidic acid ampicillin amide.

- 5 To a suspension of D-(-)-(α -aminobenzyl)penicillin
(ampicillin) (0.10g, 0.29 mmol) in 10 ml acetonitrile was
added a solution of 3-thiazolidine-2-thione-elaidylamide
and DBU (0.043ml, 0.29 mmol), and the two-phase reaction
10 mixture was stirred vigorously at ambient temperature for
72 hours. The solvents were evaporated off, and the
residue was partitioned between ethylacetate and
saturated sodium chloride(aq). The crude product
separated out, and was re-dissolved in methanol. The
mixture was evaporated to dryness to give 0.1g (55%) of
15 the title compound.

^1H NMR (DMSO- d_6 , 300 MHz) δ : 8.95(1H, d, NH), 8.5(1H, d,
NH), 7.5-7.2(5H, m, ArH), 5.75 and 5.40(2H, m, N-CHCH-S),
5.35(2H, m, CH=CH), 3.85(1H, s, CHCOO), 2.25(2H, t,
CH₂CON), 1.95(4H, m, CH₂-C=), 1.55(2H, m, CH₂-C-OOC), 1.52
20 and 1.42(6H, s, CH₃), 1.25(20H, m, CH₂), 0.85(3H, t, CH₃).

25

30

35

EXAMPLE 30

Ampicillin-oley ester

5 To a suspension of ampicillin (1.21g, 3.5 mmol) and
potassium bicarbonate (0.35g, 3.5 mmol) in 30 ml DMF was
added benzaldehyde (0.92g, 8.7 mmol) and the reaction
mixture was stirred at 0°C for 4 hours. Potassium
bicarbonate (0.35g, 3.5 mmol) and oleylbromide (1.21g,
10 3.7 mmol) was added and stirring continued at 0°C for 2
hours and at ambient temperature for 12 hours. The
solvent was evaporated at high vacuum, and the residue
was extracted with ethylacetate and cold (0°C) water. The
organic phase was evaporated to give 2.46g of a yellow
15 syrup.

The crude product was dissolved in acetonitrile and 1M
HCl(aq) was added until pH =2. 30 ml water was added and
the acetonitrile was evaporated off. The product was
extracted with ethylacetate, and dichloromethane. The
20 combined organic phases were evaporated and the crude
product was purified on a column of silica gel with 1%
triethylamine in ethylacetate as the solvent system.
Homogenous fractions were evaporated to give 0.8g (38%)
of the title compound.

25 ¹H NMR (DMSO-d₆, 300 MHz) δ: 7.5-7.2 (5H, m, ArH), 5.60
and 5.50 (2H, m, N-CHCH-S), 5.35 (2H, m, CH=CH), 4.5 (1H, m,
CH-N), 4.35 (1H, s, CH-COO), 4.05 (2H, t, CH₂-OCO),
1.95 (4H, m, CH₂-C=), 1.55 (2H, m, CH₂-C-OOC), 1.52 and
1.42 (6H, s, CH₃), 1.25 (22H, m, CH₂), 0.85 (3H, t, CH₃).

30

Antiparasitic Drugs


Parasitic infections constitute a significant problem in human and veterinary medicine. The parasites normally enter the host organism via food/water or through insect bites. The parasites may be found both in the intestine tract (epithelium cell layer) or in the blood stream where either the red blood cells or other target organs such as lungs or brain may be infected. The parasites may be found both inter- and intra-cellular in the host. Often, as for protozoas, there are stages in the life cycle of the parasites, but not all stages are subject to treatment.

The most predominant parasitic infection in man is malaria, whereas in animals, especially for birds (poultry), the intestinal infection Coccidiosis constitutes a major problem. In an untreated situation, the faeces will contain spores that will lead to re-infection of the animal or of new individuals (crossover to other species as well).

It is important to achieve an efficient transport of the active drug into the parasite itself, or as for instance with malaria and coccidiosis, into the parasite-infected cells.

The lipophilic antiparasitic derivatives of the present invention may be prepared by the general preparative methods already described.

For example, reaction Scheme 10 shows the acylation of the anti-malarial drug hydroxy-chloroquine (XXX). The reaction is quite selective on the primary OH group.



Scheme 10.

5

Biological Effects

An experiment which illustrates the enhanced anti-malarial effect of a selected hydroxychloroquine derivative is described below.

Effect of hydroxychloroquine-elaidic acid ester on malaria in mice

15

A 4-day test in NK65 strain drug sensitive *P. berghei* was carried out in Swiss albino female mice with doses of 0.063, 0.25, 1.0 and 4 mg/kg hydroxychloroquine-elaidate and 0.094, 0.395, 1.5 and 6 mg/kg hydroxychloroquine given intraperitoneally for 4 days to groups of 3 mice per group. The parasite inoculum of 10^7 infected cells was given intravenously on day 0 and followed by the drug dose for that day. The animals were dosed for the 3 following days and on the 5th day tail blood films were prepared for parasitaemia estimation. On a molar basis, the elaidic acid derivative is 2.5- 3 times more effective than hydroxychloroquine itself in the 4 day test. These findings could be of great importance in the treatment of malaria in man.

30


Compound	ED50 value (SE) mg/kg	ED90 value mg/kg	ED99 value mg/kg
Hydroxychloroquine sulphate	1.74 (1.34 - 2.14)	2.75	4.52
Hydroxychloroquine-elaidate (values x 433/600)	0.71 (0.58 - 0.83)	0.93	1.24

Table 3


- 5 The effective doses giving 50%, 90% and 99% reduction in *P. berghei* malaria parasites in the blood of mice with either hydroxychloroquine sulphate or hydroxychloroquine-elaidate is shown in this table.
- 10 Examples of antiparasitic drugs which can be derivatised in accordance with the present invention include:




15 AMODIAQUINE




HYDROXYCHLOROQUINE




20 MEFLOQUINE



MEPACRINE




PARVAQUONE




BUPARVAQUONE

5



DECOQUINATE

10



ZOALENE

15 Malaria remains the most widespread parasitic disease, with the estimated incidence of malaria in the order of 200-500 million clinical cases each year. Acquired resistance to anti-malarial drugs is a growing problem.

20 Man is naturally infected with sporozoites injected by the bite of infected female anopheline mosquitoes. The parasites rapidly leave the circulation and localize in hepatic parenchymal cells where they multiply and develop into tissue shizonts. Agents can be used for causal prophylaxis if they act on tissue forms of plasmodia within the liver. The preference for hepatic tissue of the lipid-derivatives of the present invention make the compounds more active against the tissue forms of 25 malarial disease, and might eradicate liver forms of *P. vivax* and *P. ovale*, which have some tissue parasites that persist and proliferate only later to produce relapses of erythrocytic infection months to years after 30 the primary attack.

P. falciparum accounts for over 85% of the cases of human malaria. Resistant strains of *P. falciparum* do not

accumulate high enough concentrations of the drug. Ca^{2+} channel blockers can partially restore the sensitivity to drugs like chloroquine. The cross-resistance to a number of chemically unrelated drugs is similar to the multi-drug resistance as seen in neoplastic diseases.

Fatty acids may exert an anti-malarial effect on their own (Krugliak et al, Experimental Parasitology 81, 97-105, 1995). Fatty acids like oleic, elaidic, linoleic and linoleic acids inhibited parasitemic development in mice infected with *Plasmodium vinckei petteri* or with *Plasmodium yoelii nigeriensis*.

However, the intracellular concentration needed to achieve a similar effect in man necessitates an unrealistic high intake of fatty acids.

Again, based on an embodiment of this invention, anti-malarial drug derivatives are surprisingly efficiently delivered to intracellular parasites in high concentration and can even circumvent the drug resistance mechanisms.

An example illustrating the preparation of an antimalarial derivative of this invention now follows:

30

35

EXAMPLE 31


7-chloro-4-[4-[ethyl(2-elaidoyloxyethyl)amino]1-methylbutylamino]-quinoline

5


To a solution of 7-chloro-4-[4-[ethyl(2-hydroxyethyl)amino]1-methylbutylamino]quinoline (hydroxychloroquine) (3.17g, 9.4 mmol) in 30 ml dichloromethane was added elaidic acid chloride (2.82g, 9.4 mmol) and the reaction mixture was stirred at ambient temperature for 48 hours. A small amount of methanol was added, and the solvents were evaporated at high vacuum. The residue was repeatedly purified on a column of silica gel (first run: chloroform/methanol 9:1, second run: 10 chloroform/methanol 95:5). Homogenous fractions were evaporated to give 1.18g (21%) of the title compound. 15 ¹H NMR (CDCl₃, 300 MHz) δ: 8.52(1H, d, ArH), 7.95(1H, d, ArH), 7.75(1H, d, ArH), 7.35(1H, dd, ArH), 6.42(1H, d, ArH), 5.35(2H, m, CH=CH), 4.15(2H, t, CH₂), 2.7(2H, t, CH₂), 2.55(2H, q, CH₂), 2.52(2H, t, CH₂), 2.25(2H, t, CH₂-COO), 1.95(4H, m, CH₂-C=), 1.6(4H, m, CH₂), 1.25(23H, m, CH₂), 1.0(3H, t, CH₃), 0.85(3H, t, CH₃).

25

Some further examples of drugs within other categories which can be derivatised in accordance with this invention will now be given.


30 CNS Drugs

CARBAMAZEPINE




PHENACEMID

35




PHENAGLYCODOL

5




SULTHIAME .

10




15 BENAPRYZINE




20 LEVODOPA


Cardiovascular Drugs




25 CAPTOPRIL




PHENYTOIN




VALPROIC ACID




BIPERIDEN




ENALAPRIL




5 BUNITROL



SELOKEN




10 LABETALOL



WARFARIN


- 15 The following general reaction schemes illustrate the preparation of derivatives of Warfarin and Seloken in accordance with this invention.

- 20 Scheme 11 shows the acylation of the anti-coagulant Warfarin (XXXI)



- 25 Scheme 11

- 30 Selective acylation of the hydroxy group of Seloken (XXXII) is complicated by the presence of the amino-function. The amino function is conveniently protected as the BOC derivate, and the OH group is transformed to an ester by the use of a fatty acid chloride. These reactions are shown in scheme 12.



Scheme 12

5

A specific example of reaction scheme 11 follows.

10

15

20

25

30

EXAMPLE 32

3-(α -acetonylbenzyl)-4-elaidoyloxycoumarin.

5


To a solution of 3-(α -acetonylbenzyl)-4-hydroxycoumarin (warfarin) (3.70g, 12 mmol) in 120 ml anhydrous dioxane and 25 ml pyridine was added elaidic acid chloride (3.60g, 12 mmol) and the reaction mixture was stirred at ambient temperature for 4 hours. The solvents were evaporated at high vacuum. The residue was partitioned between ether and water. The organic phase was washed with tartaric acid (aq), NaHCO₃ (aq) and water. The dried organic phase was concentrated and the product purified on a column of silica gel with heptane/ethylacetate (6:1) as the eluent system. Impure fractions were repurified and homogenous fractions were evaporated to give 5.1g (70%) of the title compound as a pale yellow oil.

¹H NMR (CDCl₃, 300 MHz) δ : 7.55- 7.15(9H, m, ArH), 5.38(2H, m, CH=CH), 4.78(1H, t, CH), 3.45(2H, m, CH₂-COCH), 2.75(2H, t, CH₂-COO), 2.18(3H, s, CH₃), 1.95(4H, m, CH₂-CH=); 1.85(2H, m, CH₂-C-COO), 1.5 - 1.2(20H, m, CH₂), 0.85(3H, t, CH₃).

As previously stated, the present invention is generally applicable to all classes of biologically active compounds and not just to compounds useful in human or animal medicine. In particular, agrochemicals which have one or more functional groups selected from alcohol, ether, phenyl, amino, amido, thiol, carboxylic acid and carboxylic acid ester groups can be derivatised in accordance with this invention. Examples of such agricultural and horticultural chemicals include:


35

93




AMINOTRIAZOLE


5




10 BENAZOLIN



ASULAM




BROMOFENOXIM




15

BROMOXYNIL




2, 4 - D



20


DICAMBA




DICLOBUTRAZOL

25

94




DINOTERB




FLUAZIFOP

5




MECOPROP




PICLORAM

10




SULFOMETHURON




20

METHAMIDOPHOS




TRICHLOROPHON




25

ANCYMIDOL




HORMODIN

95




CYCLOHEXIMIDE



HYMEXAZOL

5



ETHIRIMOL

10

15

20

25

30

CLAIMS:

- 5 1. A lipophilic derivative of a biologically
active compound containing in its molecular structure one
or more functional groups selected from alcohol, ether,
phenyl, amino, amido, thiol, carboxylic acid and
carboxylic acid ester groups, other than a nucleoside or
10 nucleoside derivative, said lipophilic derivative being
characterised by a molecular structure in which the or at
least one said functional group of said biologically
active compound is replaced by a lipophilic group
selected from those of the formula: RCOO- , RCONH- ,
15 RCOS- , $\text{RCH}_2\text{O-}$, $\text{RCH}_2\text{NH-}$, $\text{-COOCH}_2\text{R}$, $\text{-CONHCH}_2\text{R}$ and $\text{-SCH}_2\text{R}$,
wherein R is a lipophilic moiety selected from cis-8-
heptadecenyl, trans-8-heptadecenyl, cis-10-nonadecenyl
and trans-10-nonadecenyl.
- 20 2. A lipophilic derivative according to Claim 1,
wherein said biologically active compound is a compound
having therapeutic activity in human or animal medicine.
3. A lipophilic derivative according to Claim 2,
25 wherein said biologically active compound is an
adrenocorticosteroid.
4. A lipophilic derivative according to Claim 3,
wherein said adrenocorticosteroid is selected from
30 betamethasone, cortisone, dexamethasone, fluocinolone,
fludrocortisone, hydrocortisone, methylprednisolone,
prednisolone, triamcinolone, eprozinol, paramethasone,
prednisone, beclomethasone and orciprenaline.
- 35 5. A lipophilic derivative according to Claim 4,
wherein said adrenocorticosteroid is selected from
prednisolone and betamethasone.

6. The compound: 11 β ,17 α ,21-trihydroxy-pregna-1,4-diene-3,20-dione-21-elaidate.
7. The compound: 9-fluoro-11 β ,17,21-trihydroxy-
5 16 β -methylpregna-1,4-diene-3,20-dione-21-elaidate.
8. A lipophilic derivative according to Claim 2, wherein said biologically active compound is a non-steroidal antiinflammatory drug (NSAID).
- 10 9. A lipophilic derivative according to Claim 8, wherein said NSAID is selected from acemetacin, alclofenac, amfenac, aspirin, bendazac, benorylate, benoxaprofen, bucloxic acid, bufexamac, bumadizon,
15 butibufen, carprofen, cinmetacin, clidanac, clometacin, cloripac, diclofenac, diflunisal, etodolac, etofenamate, felbinac, fenbufen, fenclofenac, fenclorac, fendosal, fenoprofen, fentiazac, flufenamic acid, flurbiprofen, glafenine, ibufenac, ibuprofen, indomethacin, isofezolac,
20 isoxepac, ketoprofen, ketorolac, lonazolac, meclofenamic acid, mefanamic acid, metiazinic acid, nabumetone, naproxen, niflumic acid, oxametacin, oxaprozin, pirazolac, piroxicam, protizinic acid, salicylic acid, sulindac, surgam, tenidap, tenoxicam, tiaramide,
25 tinoridine, tolfenamic acid, tolmetin and zomepirac.
10. A lipophilic derivative according to Claim 9, wherein said NSAID is selected from diclofenac, indomethacin, naproxen and piroxicam.
- 30 11. The compound: (2-[2,6-dichlorophenyl]amino-benzeneacetic acid)-(cis-9'-octadecenyl)ester.
12. The compound: 1-(p-chlorobenzoyl)-5-methoxy-
35 2-methylindole-3-acetic acid (cis-9-octadecenyl)amide.

13. The compound: S(+)-2-(6-methoxy-2-naphthyl)-propionic acid (cis-9'-octadecenyl)amide.
- 5 14. The compound: S(+)-2-(6-methoxy-2-naphthyl)-propionic acid-cis-9'-octadecenyl ester.
15. The compound: 4-O-(trans-9'-octadecenoyl)-2-methyl-N(2-pyridyl)-2H-1,2-benzothiazine-3-carboxamide-1,1-dioxide.
- 10 16. The compound: 4-O-(cis-11'-eicosenoyl)-2-methyl-N(2-pyridyl)-2H-1,2-benzothiazine-3-carboximide-1,1-dioxide.
- 15 17. The compound: 2-hydroxy-benzoic acid-(cis-9'-octadecenyl)ester.
18. The compound: 2-(cis-9'-octadecenoxyl)-ethyl benzoate.
- 20 19. The compound: 2-(cis-9'-octadecenoxyl)-benzoic acid.
- 25 20. The compound: S(+)-2-(6-[cis-9'-octadecenoxyl]-2-naphthyl)-propionic acid ethyl ester.
21. The compound: S(+)-2-(6-[cis-9'-octadecenoxyl]-2-naphthyl)-propionic acid.
- 30 22. A lipophilic derivative according to Claim 2, wherein said biologically active compound is an anti-cancer drug.

23. A lipophilic derivative according to Claim 22,
wherein said anti-cancer drug is selected from megestrol,
medroxyprogesterone, hexestrol, trilostane, amino-
5 glutethimide, epitio stanol, calusterone, podophyllinic
acid 2-ethylhydrazide, pirarubicin, doxorubicin,
daunorubicin, taxol, mopidamol, mitoxantrone, lonidamine,
etoposide, eflornitine, defosamide, trimetrexate,
methotrexate, deopterin, thioguanin, thiamiprene,
10 mercaptopurin, dacarbazine, nimustine, chlorozotocin,
melphalan, estramustin, cyclophosphamide, chlorambucil
and trimethylmelamine.

24. A lipophilic derivative according to Claim 23,
15 wherein said anti-cancer drug is selected from
chlorambucil, melphalan and taxol.

25. The compound: chlorambucil-oleyl ester.

20 26. The compound: elaidic acid mephalan amide.

27. The compound: taxol-2'-elaidate.

28. The compound: elaidic acid daunorubicin
25 amide.

29. The compound: elaidic acid doxorubicin amide.


30 30. The compound: daunorubicine oleyl carbamate.

31. The compound: doxorubicin oleyl carbamate.


32. A lipophilic derivative according to Claim 2,
wherein said biologically active compound is an
35 antimicrobial agent.

33. A lipophilic derivative according to Claim 32,
wherein said antimicrobial agent is selected from
oxacillin, ampicillin, amoxicillin, cephalixin,
cephalotin, cephalosporin, doxycyclin, chloramphenicol,
5 p-amino-salicylic acid, ethambutol, ciprofloxacin,
enrofloxacin, difloxacin and danofloxacin.
34. A lipophilic derivative according to Claim 33,
wherein said antimicrobial agent is selected from p-amino
10 salicylic acid, chloramphenicol, oxacillin and
ampicillin.
35. The compound: p-amino-salicylic acid
elaidylester.
15
36. The compound: 4-(elaidamido)salicylic acid.
37. The compound: elaidic acid chloramphenicol
ester.
20
38. The compound: oxacillin oleyl ester.
39. The compound: elaidic acid ampicillin amide.
- 25 40. The compound: ampicillin oleyl ester.
41. A lipophilic derivative according to Claim 2,
wherein said biologically active compound is an
antiparasitic drug.
30

42. A lipophilic derivative according to Claim 41,
wherein said antiparasitic drug is selected from
amodiaquine, hydroxychloroquine, mefloquine, mepacrine,
5 decoquinate, zoalene, a compound of the formula:



and a compound of the formula:



10

43. The compound: 7-chloro-4-[4-[ethyl(2-
elaidoyloxyethyl)-amino]]-methylbutylamino]quinolone.
44. A lipophilic derivative according to Claim 2,
15 wherein said biologically active compound is a CNS drug.
45. A lipophilic derivative according to Claim 44,
wherein said CNS drug is selected from carbamazepine,
phenacemid, phenaglycodol, phenytoin, sulthiame, valproic .
20 acid, benapryazine, biperiden and levodopa.
46. A lipophilic derivative according to Claim 2,
wherein said biologically active compound is a
cardiovascular drug.
- 25 47. A lipophilic derivative according to Claim 46,
wherein said cardiovascular drug is selected from
captopril, enalapril, bunitrol, seloken, labetalol and
seloken.

48. The compound: 3-(α -acetylbenzyl)-4-elaidoyloxy coumarin.

49. A pharmaceutical preparation, comprising a
5 compound according to any one of Claims 2-47 and a
pharmaceutically acceptable carrier or excipient.

50. A lipophilic derivative according to Claim 1,
wherein said biologically active compound is an
10 agrochemical.

51. A lipophilic derivative according to Claim 50,
wherein said agrochemical is pesticide, fungicide or
herbicide.

15 52. A lipophilic derivative according to Claim 50,
wherein said agrochemical is selected from aminotriazole,
asulam, benazolin, bromofenoxim, bromoxynil, 2,4-D,
DICAMBA, diclobutrazol, dinoterb, fluazifop, mecoprop,
20 picloram, sulfomethuron, methamidophos, trichlorophon,
ancymidol, hormodin, cycloheximide, hymexazol and
ethirimol.

53. A composition useful in agriculture and
25 horticulture, comprising a compound according to any one
of Claims 50-52 and a carrier or diluent therefor.

30

35

54. A method for modifying the activity of a biologically active compound containing in its molecular structure one or more functional groups selected from
5 alcohol, ether, phenyl, amino, amido, thiol, carboxylic acid and carboxylic acid ester groups, other than a nucleoside or nucleoside analogue, comprising replacing the or at least one said functional group of said biologically active compound by a lipophilic group
10 selected from those of the formula: RCOO- , RCONH- , RCOS- , $\text{RCH}_2\text{O-}$, $\text{RCH}_2\text{NH-}$, $\text{-COOCH}_2\text{R}$, $\text{-CONHCH}_2\text{R}$ and $\text{-SCH}_2\text{R}$, wherein R is a lipophilic moiety selected from cis-8-heptadecenyl, trans-8-heptadecenyl, cis-10-nonadecenyl and trans-10-nonadecenyl.

15

55. A method of preparing a lipophilic derivative according to Claim 1, characterised in that said biologically active compound is reacted with a cis or trans-n-9 monounsaturated fatty acid, fatty alcohol or
20 fatty amine having a chain length of 18 or 20 carbon atoms, or with a reactive derivative of such fatty acid, fatty alcohol or fatty amine.


25

30

35


1/13

Fig. 1.




2/13

Fig. 2.




3/13

Fig. 3.




4/13

Fig. 4.




5/13

Fig. 5.




6/13

Fig. 6.




7/13

Fig. 7.




8/13

Fig. 8.



9/13


Fig. 9.



10/13


Fig. 10.

Bronchial responsiveness to 5HT




11/13

Fig. 11.



12/13


Fig. 12.



- ◆— Daunorubicine-elaidic-amide CEM
- Daunorubicine-elaidate-amide VLB
- ▲— Daunorubicine-oleic-carbamate CEM
- ×— Daunorubicine-oleyl-carbamate VLB

13/13

Fig. 13.



1
INTERNATIONAL SEARCH REPORT

International application No.
PCT/NO 98/00021

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07C:, C07D:, C07J:, A61K:, A01N:
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6:

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CA, WPI

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0105404 A1 (MERCK & CO. INC.), 18 April 1984 (18.04.84) --	1-7, 49, 54-55
P, X	EP 0771817 A2 (LABORATORIOS S.A.L.V.A.T., S.A.), 7 May 1997 (07.05.97) --	1-7, 49, 54-55
X	International Journal of Obesity, Volume 20, 1996, D Sanchis et al, "Oleoyl-estrone induces the loss of body fat in rats" --	1-7, 49, 54-55
X	EP 0245687 A1 (WARNER-LAMBERT COMPANY), 19 November 1987 (19.11.87) --	1-2, 49, 54-55

☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.

* Special categories of cited documents	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document but published on or after the international filing date	"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 23 June 1998	Date of mailing of the international search report 25 -06- 1998
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86	Authorized officer Eva Johansson Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/NO 98/00021

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0054435 A1 (FUJISAWA PHARMACEUTICAL CO. LTD.), 23 June 1982 (23.06.82) --	1-2,49,54-55
A	WO 9525504 A1 (PHARMAVENE, INC.), 28 Sept 1995 (28.09.95) --	1-2,49,54-55
A	US 5447729 A (GEORGE W. BELENDUIK ET AL), 5 Sept 1995 (05.09.95) --	1-2,49,54-55
X	WO 9001477 A1 (GÖDEKKE AKTIENGESELLSCHAFT), 22 February 1990 (22.02.90) --	1-2,8,22,44, 46,49,54-55
X	EP 0255126 A2 (GÖDECKE AKTIENGESELLSCHAFT), 3 February 1988 (03.02.88) --	1-2,22,44, 46,49,54-55
X	EP 0508690 A1 (AMERICAN HOME PRODUCTS CORPORATION), 14 October 1992 (14.10.92) --	1-2,8,49, 54-55
X	EP 0148752 A2 (G.D. SEARLE & CO.), 17 July 1985 (17.07.85) --	1-2,8,49, 54-55
X	EP 0282127 A2 (THE PROCTER & GAMBLE COMPANY), 14 Sept 1988 (14.09.88) --	1-2,8,49, 54-55
X	EP 0579066 A2 (F. HOFFMANN-LA ROCHE AG), 19 January 1994 (19.01.94) --	1-2,8,49, 54-55
X	EP 0206609 A1 (THE PROCTER & GAMBLE COMPANY), 30 December 1986 (30.12.86) --	1-2,8,49, 54-55

INTERNATIONAL SEARCH REPORT

International application No.

PCT/NO 98/00021

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0359347 A2 (NEORX CORPORATION), 21 March 1990 (21.03.90) --	1-2, 22-24, 49, 54-55
A	WO 9320800 A1 (KINNUNEN, PAAVO, KAI, JOHANNES), 28 October 1993 (28.10.93) --	1-2, 22, 49, 54-55
X	WO 9426262 A1 (PUBLIC HEALTH RESEARCH INSTITUTE), 24 November 1994 (24.11.94) --	1-2, 32-40, 49, 54-55
A	US 3882243 A (TADAO MAEDA ET AL), 6 May 1975 (06.05.75) --	1-2, 32-34, 37, 49, 54-55
A	FR 2485001 A (SVATOS ANTONIN), 24 December 1981 (24.12.81) --	1-2, 41, 49, 54-55
A	US 4554279 A (ANDREW J. SAGGIOMO ET AL), 19 November 1985 (19.11.85) --	1-2, 41, 49, 54-55
X	US 5284876 A (VICTOR E. SHASHOUA ET AL), 8 February 1994 (08.02.94) --	1-2, 46, 49, 54-55
A	WO 9308841 A1 (SRI INTERNATIONAL), 13 May 1993 (13.05.93) --	1-2, 46-47, 49, 54-55
A	WO 9530415 A1 (EMORY UNIVERSITY), 16 November 1995 (16.11.95) --	1-2, 46-47, 49, 54-55

INTERNATIONAL SEARCH REPORT

International application No.

PCT/NO 98/00021

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>STN International, File CAPLUS, CAPLUS accession no. 120:263749, Serre, I. et al: "Seed oil derivatives as adjuvants: influence of methyl to octadecyl oleates on the penetration of herbicides through various plant cuticles"; & Meded. - Fac. Landbouwk. Toegepaste Biol. Wet. (Univ. Gent) (1993), 58(3a), 795-802</p> <p style="text-align: center;">-- -----</p>	<p>1-2,50-51, 53,54-55</p>

INTERNATIONAL SEARCH REPORT

International application No.

PCT/NO 98/00021

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(e).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see next sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☒

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/NO 98/00021

Lack of unity a posteriori

The general problem underlying the invention is not novel and a solution to it has already been found or does not involve an inventive step having regard to the state as illustrated by:

- a) International Journal of Obesity (1996) 20, pages 588-594
- b) EP 0105404 A1
- c) EP 0579066 A2
- d) EP 0282127 A2
- e) EP 0206609 A1
- f) EP 0148752 A2
- g) WO 90/01477
- h) EP 0255126 A2
- i) US 5284876
- j) EP 0054435

Therefore, the original single general inventive concept is not acceptable anymore, making it necessary to reconsider the technical relationship between the different solutions mentioned. This leads to their regrouping under distinct subjects as listed below, each subject now falling under its own inventive concept.

- 1. Claims 1-7, 49, 54-55: Concerning steroids
- 2. Claims 1-2, 8-21, 49, 54-55: Concerning non-steroidal anti-inflammatory drugs
- 3. Claims 1-2, 22-31, 49, 54-55: Concerning anti-cancer agents
- 4. Claims 1-2, 32-40, 49, 54-55: Concerning anti-microbial agents
- 5. Claims 1-2, 41-43, 49, 54-55: Concerning anti-parasitic agents
- 6. Claims 1-2, 44-45, 49, 54-55: Concerning CNS drugs
- 7. Claims 1-2, 46-48, 49, 54-55: Concerning cardiovascular drugs
- 8. Claims 1-2, 50-53, 54-55: Concerning agrochemical compounds

INTERNATIONAL SEARCH REPORT
Information on patent family members

09/06/98

International application No.

PCT/NO 98/00021

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0105404 A1	18/04/84	JP 59082399 A US 4560512 A	12/05/84 24/12/85
EP 0771817 A2	07/05/97	ES 1031812 U,Y	16/02/96
EP 0245687 A1	19/11/87	SE 0245687 T3 CA 1264747 A JP 62258366 A US 4722927 A	23/01/90 10/11/87 02/02/88
EP 0054435 A1	23/06/82	SE 0054435 T3 AT 5961 E,T CA 1170599 A JP 1663251 C JP 3030586 B JP 57131747 A US 4436726 A	15/02/84 10/07/84 19/05/92 30/04/91 14/08/82 13/03/84
WO 9525504 A1	28/09/95	AU 2102495 A CA 2185803 A EP 0788346 A JP 9510712 T	09/10/95 28/09/95 13/08/97 28/10/97
US 5447729 A	05/09/95	AU 2276095 A CA 2187202 A EP 0754031 A JP 9511744 T WO 9527479 A	30/10/95 19/10/95 22/01/97 25/11/97 19/10/95
WO 9001477 A1	22/02/90	AU 627766 B AU 4048989 A CN 1040193 A DE 3826846 A DK 11391 A EP 0354495 A EP 0427765 A FI 910554 D JP 4500958 T NO 177785 B,C PT 91351 A,B US 5185352 A US 5322844 A	03/09/92 05/03/90 07/03/90 08/02/90 23/01/91 14/02/90 22/05/91 00/00/00 20/02/92 14/08/95 08/03/90 09/02/93 21/06/94

INTERNATIONAL SEARCH REPORT
Information on patent family members

09/06/98

International application No.

PCT/NO 98/00021

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0255126 A2	03/02/88	SE 0255126 T3	
		AU 597090 B	24/05/90
		AU 7624887 A	04/02/88
		CA 1318670 A	01/06/93
		DE 3625738 A	11/02/88
		DE 3782207 A	19/11/92
		DK 374887 A	31/01/88
		ES 2052525 T	16/07/94
		FI 89042 B	30/04/93
		FI 873313 A	31/01/88
		JP 63048250 A	29/02/88
		SU 1470177 A	30/03/89
		US 4855489 A	08/08/89
EP 0508690 A1	14/10/92	AU 649120 B	12/05/94
		AU 1482692 A	15/10/92
		CA 2065228 A	13/10/92
		FI 921596 A	13/10/92
		IL 101444 A	18/02/97
		JP 5097839 A	20/04/93
		MX 9201657 A	01/10/92
		NZ 242301 A	24/03/97
		US 5242945 A	07/09/93
		US 5679801 A	21/10/97
		ZA 9202181 A	27/09/93
EP 0148752 A2	17/07/85	EP 0305958 A	08/03/89
		JP 60208951 A	21/10/85
		US 4551279 A	05/11/85
EP 0282127 A2	14/09/88	SE 0282127 T3	
		AU 621041 B	05/03/92
		AU 1279588 A	27/10/88
		CA 1328457 A	12/04/94
		DE 3887544 D,T	01/06/94
		DK 127588 A	10/09/88
		EG 18362 A	30/10/92
		ES 2061624 T	16/12/94
		FI 98812 B,C	15/05/97
		FI 881064 A	10/09/88
		HK 94596 A	07/06/96
		IE 60992 B	07/09/94
		JP 2562475 B	11/12/96
		JP 63295537 A	01/12/88
		KR 9612204 B	16/09/96
		NO 180117 B,C	11/11/96
		PT 86918 B	29/05/92
		US 5045565 A	03/09/91
		US 5099030 A	24/03/92
		ZA 8801677 A	06/09/88

INTERNATIONAL SEARCH REPORT
Information on patent family members

09/06/98

International application No.

PCT/NO 98/00021

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5284876 A	08/02/94	AU 635203 B	18/03/93
		AU 3354689 A	22/09/89
		DK 526189 A	12/12/89
		EP 0401301 A	12/12/90
		FI 904207 D	00/00/00
		JP 3503893 T	29/08/91
		US 4933324 A	12/06/90
		US 4939174 A	03/07/90
		WO 8907938 A	08/09/89
WO 9308841 A1	13/05/93	US 5238933 A	24/08/93
WO 9530415 A1	16/11/95	AU 2586095 A	29/11/95
		BG 101030 A	30/09/97
		BR 9507716 A	23/09/97
		CA 2189336 A	16/11/95
		EP 0759752 A	05/03/97
		HU 76728 A	28/11/97
		HU 9603041 D	00/00/00
		JP 10500111 T	06/01/98
		NO 964742 A	08/11/96
		PL 317193 A	17/03/97
		US 5750351 A	12/05/98